

Anti-CD28 Antibody

Catalog # AP95546

Product Information

Application	WB, FC, IF/IC
Primary Accession	P10747
Reactivity	Human
Host	Mouse
Clonality	Monoclonal (clone: CD28.2)
Calculated MW	25066


Additional Information

Gene ID	940
Other Names	T-cell-specific surface glycoprotein CD28; TP44; CD28
Target/Specificity	Native purified human CD28
Dilution	WB~~WB (1/500 - 1/1000) FC~~1:10~50 IF/IC~~N/A
Format	Mouse IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CD28
Function	Receptor that plays a role in T-cell activation, proliferation, survival and the maintenance of immune homeostasis (PubMed:1650475, PubMed:7568038). Functions not only as an amplifier of TCR signals but delivers unique signals that control intracellular biochemical events that alter the gene expression program of T-cells (PubMed:24665965). Stimulation upon engagement of its cognate ligands CD80 or CD86 increases proliferation and expression of various cytokines in particular IL2 production in both CD4(+) and CD8(+) T-cell subsets (PubMed:12196291, PubMed:1650475, PubMed:35397202). Mechanistically, ligation induces recruitment of protein kinase C-theta/PRKCQ and GRB2 leading to NF-kappa-B activation via both PI3K/Akt-dependent and -independent pathways (PubMed:21964608, PubMed:24665965, PubMed:7568038). In conjunction with TCR/CD3 ligation and CD40L costimulation, enhances the production of IL4 and IL10 in T- cells (PubMed:8617933).
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Expressed in T-cells and plasma cells, but not in less mature B-cells

Images



Western blot analysis of CD28 expression in U87 (A), HuH7 (B), HuccT1 (C) whole cell lysates. (Predicted band size: 25 kD; Observed band size: 37 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.