

WIPI2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9559b

Product Information

Application	WB, E
Primary Accession	Q9Y4P8
Other Accession	Q6AY57 , Q80W47 , Q5ZHN3
Reactivity	Human, Mouse
Predicted	Chicken, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB24217
Calculated MW	49408
Antigen Region	426-454

Additional Information

Gene ID	26100
Other Names	WD repeat domain phosphoinositide-interacting protein 2, WIPI-2, WIPI49-like protein 2, WIPI2
Target/Specificity	This WIPI2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 426-454 amino acids from the C-terminal region of human WIPI2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WIPI2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	WIPI2 (HGNC:32225)
Function	Component of the autophagy machinery that controls the major intracellular degradation process by which cytoplasmic materials are

packaged into autophagosomes and delivered to lysosomes for degradation (PubMed:[20505359](#), PubMed:[28561066](#)). Involved in an early step of the formation of preautophagosomal structures (PubMed:[20505359](#), PubMed:[28561066](#)). Binds and is activated by phosphatidylinositol 3-phosphate (PtdIns3P) forming on membranes of the endoplasmic reticulum upon activation of the upstream ULK1 and PI3 kinases (PubMed:[28561066](#)). Mediates ER-isolation membranes contacts by interacting with the ULK1:RB1CC1 complex and PtdIns3P (PubMed:[28890335](#)). Once activated, WIPI2 recruits at phagophore assembly sites the ATG12-ATG5-ATG16L1 complex that directly controls the elongation of the nascent autophagosomal membrane (PubMed:[20505359](#), PubMed:[28561066](#)).

Cellular Location

Preautophagosomal structure membrane; Peripheral membrane protein; Cytoplasmic side. Note=Localizes to omegasomes membranes which are endoplasmic reticulum connected structures at the origin of preautophagosomal structures. Enriched at preautophagosomal structure membranes in response to PtdIns3P.

Tissue Location

Ubiquitously expressed (at protein level). Highly expressed in heart, skeletal muscle and pancreas. Expression is down-regulated in pancreatic and in kidney tumors


Background

WD40 repeat proteins are key components of many essential biologic functions. They regulate the assembly of multiprotein complexes by presenting a beta-propeller platform for simultaneous and reversible protein-protein interactions. Members of the WIPI subfamily of WD40 repeat proteins, such as WIPI2, have a 7-bladed propeller structure and contain a conserved motif for interaction with phospholipids (Proikas-Cezanne et al., 2004 [PubMed 15602573]).

References

Proikas-Cezanne, T., et al. Oncogene 23(58):9314-9325(2004)
Simpson, J.C., et al. EMBO Rep. 1(3):287-292(2000)

Images



Western blot analysis of WIPI2 Antibody (C-term) (Cat. #AP9559b) in mouse brain tissue lysates (35ug/lane). WIPI2 (arrow) was detected using the purified Pab.

Citations

- [Automated Detection of Autophagy Response Using Single Cell-Based Microscopy Assays.](#)
- [WIPI3 and WIPI4 \$\beta\$ -propellers are scaffolds for LKB1-AMPK-TSC signalling circuits in the control of autophagy.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.