

Anti-APC10 Antibody

Catalog # AP95669

Product Information


Application	WB, IHC, FC, IF/IC
Primary Accession	Q9UM13
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	21252

Additional Information


Gene ID	10393
Other Names	APC10; Anaphase-promoting complex subunit 10; APC10; Cyclosome subunit 10
Target/Specificity	Recombinant fusion protein of human APC10 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200) IF/IC~~N/A
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	ANAPC10
Synonyms	APC10
Function	Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through the cell cycle (PubMed:18485873, PubMed:27120157, PubMed:27509861). APC/C acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (PubMed:18485873). APC/C catalyzes assembly of branched 'Lys-11'-/'Lys- 48'-linked branched ubiquitin chains on target proteins (PubMed:29033132). APC/C is activated by CDC20 in the metaphase/anaphase transition of cell cycle, targeting the degradation of cyclin B and securin (PubMed:27120157, PubMed:27509861). APC/C is regulated by the mitotic checkpoint complex (MCC), which inhibits APC/C and delays chromosome segregation (PubMed:27509861).




Western blot analysis of APC10 expression in HeLa (A), MCF7 (B), SKBr3 (C), A431 (D), HEK293 (E), A549 (F), SPCA1 (G) whole cell lysates. (Predicted band size: 21 kDa; Observed band size: 23 kDa)



Immunohistochemical analysis of APC10 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of APC10 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



Flow cytometric analysis of HeLa cells using Anti-APC10 Antibody (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.