

Anti-Caveolin 2 Antibody

Catalog # AP95719

Product Information

Application	WB, IHC, FC
Primary Accession	P51636
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	18291

Additional Information

Gene ID	858
Other Names	Caveolin-2
Target/Specificity	Recombinant fusion protein of human Caveolin 2 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200)
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	CAV2
Function	May act as a scaffolding protein within caveolar membranes. Interacts directly with G protein alpha subunits and can functionally regulate their activity. Acts as an accessory protein in conjunction with CAV1 in targeting to lipid rafts and driving caveolae formation. The Ser-36 phosphorylated form has a role in modulating mitosis in endothelial cells. Positive regulator of cellular mitogenesis of the MAPK signaling pathway. Required for the insulin-stimulated nuclear translocation and activation of MAPK1 and STAT3, and the subsequent regulation of cell cycle progression (By similarity).
Cellular Location	Nucleus. Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Cell membrane; Peripheral membrane protein. Membrane, caveola; Peripheral membrane protein. Note=Potential hairpin-like structure in the membrane. Membrane protein of caveolae Tyr-19-phosphorylated form is enriched at sites of cell-cell contact and is translocated to the nucleus in complex with MAPK1 in response to insulin (By similarity). Tyr-27-phosphorylated form is located both in the cytoplasm and plasma membrane. CAV1-mediated Ser-23-phosphorylated form locates to the plasma membrane. Ser-36-phosphorylated form resides in intracellular

compartments.


Tissue Location

Expressed in endothelial cells, smooth muscle cells, skeletal myoblasts and fibroblasts


Images



Western blot analysis of Caveolin 2 expression in A549 (A), 3T3L1 (B), A431 (C) whole cell lysates. (Predicted band size: 18 kDa; Observed band size: 18 kDa)



Immunohistochemical analysis of Caveolin 2 staining in human endometrial cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Flow cytometric analysis of A549 cells using Anti-Caveolin 2 Antibody (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.