

Anti-CD14 Antibody

Catalog # AP95729

Product Information

Application	WB, IHC, FC
Primary Accession	P08571
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Calculated MW	40076

Additional Information

Gene ID	929
Other Names	Monocyte differentiation antigen CD14; Myeloid cell-specific, leucine-rich, h glycoprotein; CD14
Target/Specificity	Recombinant fusion protein of human CD14 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200)
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	CD14
Function	Coreceptor for bacterial lipopolysaccharide (PubMed: 1698311 , PubMed: 23264655). In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the LY96/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed: 20133493 , PubMed: 22265692 , PubMed: 23264655). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed: 8612135). Acts as a coreceptor for TLR2:TLR6 heterodimer in response to diacylated lipopeptides and for TLR2:TLR1 heterodimer in response to triacylated lipopeptides, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway (PubMed: 16880211). Binds electronegative LDL (LDL(-)) and mediates the cytokine release induced by LDL(-) (PubMed: 23880187).
Cellular Location	Cell membrane; Lipid-anchor, GPI-anchor. Secreted. Membrane raft. Golgi apparatus. Note=Secreted forms may arise by cleavage of the GPI anchor.


Tissue Location

Detected on macrophages (at protein level) (PubMed:1698311). Expressed strongly on the surface of monocytes and weakly on the surface of granulocytes; also expressed by most tissue macrophages.


Images



Western blot analysis of CD14 expression in HepG2 (A), A549 (B), HL60 (C), RAW264.7 (D), Hela (E), HEK293 (F), NIH/3T3 (G) whole cell lysates. (Predicted band size: 40 kD; Observed band size: 45 kD)



Immunohistochemical analysis of CD14 staining in human stomach cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Flow cytometric analysis of Jurkat cells using Anti-CD14 Antibody (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.