

Anti-ETFA Antibody

Catalog # AP95822

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	P13804
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	35080

Additional Information


Gene ID	2108
Other Names	Electron transfer flavoprotein subunit alpha, mitochondrial; Alpha-ETF
Target/Specificity	Recombinant fusion protein of human ETFA expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200) IF/IC~~N/A
Format	Mouse IgG2a. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	ETFA
Function	Heterodimeric electron transfer flavoprotein that accepts electrons from several mitochondrial dehydrogenases, including acyl-CoA dehydrogenases, glutaryl-CoA and sarcosine dehydrogenase (PubMed: 10356313 , PubMed: 15159392 , PubMed: 15975918 , PubMed: 27499296 , PubMed: 9334218). It transfers the electrons to the main mitochondrial respiratory chain via ETF-ubiquinone oxidoreductase (ETF dehydrogenase) (PubMed: 9334218). Required for normal mitochondrial fatty acid oxidation and normal amino acid metabolism (PubMed: 12815589 , PubMed: 1430199 , PubMed: 1882842).
Cellular Location	Mitochondrion matrix

Images


Western blot analysis of ETFA expression in HepG2 (A),




A431 (B), HEK293 (C), HeLa (D), MCF7 (E) whole cell lysates.
(Predicted band size: 35 kD; Observed band size: 35 kD)



Immunohistochemical analysis of ETFA staining in human cervical cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ETFA staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



Flow cytometric analysis of HeLa cells using Anti-ETFAs Antibody (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.