

Anti-CD223 Antibody

Catalog # AP95923

Product Information

Application	WB, FC
Primary Accession	P18627
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	57449

Additional Information

Gene ID	3902
Other Names	FDC; Lymphocyte activation gene 3 protein; LAG-3; Protein FDC; CD223
Target/Specificity	Recombinant fusion protein of human CD223 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) FC~~FC (1/100 - 1/200)
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	LAG3 {ECO:0000303 PubMed:35761082, ECO:0000312 HGNC:HGNC:6476}
Function	[Lymphocyte activation gene 3 protein]: Inhibitory receptor on antigen activated T-cells (PubMed: 20421648 , PubMed: 35761082 , PubMed: 7805750 , PubMed: 8647185). Delivers inhibitory signals upon binding to ligands, such as MHC class II, its main ligand present at the surface of antigen-presenting cells (APCs), and FGL1, which is secreted by hepatocytes and certain types of tumor cells (PubMed: 30580966 , PubMed: 32920841 , PubMed: 35761082 , PubMed: 39671469 , PubMed: 7589152 , PubMed: 8647185 , PubMed: 9159144). Ligand-binding initiates a signaling that inhibits the T-cell receptor (TCR) in the immunological synapse, preventing T-cell activation (PubMed: 40101708). Mechanistically, ligand-binding promotes (1) ubiquitination of the KIEELE motif, unleashing the RRFSALE motif from the membrane and (2) leading to the formation of condensates with the TCR component CD3E, thereby disrupting the association between CD3E and LCK and preventing TCR activation (PubMed: 40101708 , PubMed: 40592325). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed: 20421648 , PubMed: 7805750 , PubMed: 8647185). Also mediates immune tolerance: constitutively expressed

on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity).


Cellular Location

[Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein. Note=Clusters on the T-cell surface following ligand-binding


Tissue Location

Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.

Images



Western blot analysis of CD223 expression in Raji (A), Ramos (B), MOLT4 (C) whole cell lysates. (Predicted band size: 57 kD; Observed band size: 95 kD)



Flow cytometric analysis of Jurkat cells using Anti-CD223 Antibody (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.