

Anti-CD273 Antibody

Catalog # AP95995

Product Information

Application	WB, FC, IF/IC
Primary Accession	Q9BQ51
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Calculated MW	30957


Additional Information

Gene ID	80380
Other Names	B7DC; CD273; PDCD1L2; PDL2; Programmed cell death 1 ligand 2; PD-1 ligand 2; PD-L2; PDCD1 ligand 2; Programmed death ligand 2; Butyrophilin B7-DC; B7-DC; CD273
Target/Specificity	Recombinant fusion protein of human CD273 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) FC~~FC (1/100 - 1/200) IF/IC~~N/A
Format	Mouse IgG2b. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	PDCD1LG2
Synonyms	B7DC, CD273, PDCD1L2, PDL2
Function	Plays a critical role in induction and maintenance of immune tolerance to self (PubMed: 11224527 , PubMed: 12538684). Acts as a ligand for the inhibitory receptor PDCD1/PD-1, inhibiting T-cell proliferation by blocking cell cycle progression and cytokine production (PubMed: 11224527).
Cellular Location	[Isoform 3]: Secreted [Isoform 1]: Cell membrane; Single-pass type I membrane protein
Tissue Location	Highly expressed in heart, placenta, pancreas, lung and liver and weakly expressed in spleen, lymph nodes and thymus


Images



Western blot analysis of CD273 expression in HepG2 (A), HeLa (B), NIH/3T3 (C), PC12 (D), K562 (E), Jurkat (F), MCF7 (G), HEK293-6e (H) whole cell lysates. (Predicted band size: 32 kD; Observed band size: 58 kD)



Immunofluorescent analysis of CD273 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



Flow cytometric analysis of HeLa cells using Anti-CD273 Antibody (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.