

Anti-HIWI2 Antibody

Catalog # AP96002

Product Information

Application	WB, IHC, FC
Primary Accession	Q7Z3Z4
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	96589

Additional Information

Gene ID	143689
Other Names	HIWI2; PIWI; Piwi-like protein 4
Target/Specificity	Recombinant fusion protein of human HIWI2 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200)
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	PIWIL4
Synonyms	HIWI2, PIWI
Function	Plays a central role during spermatogenesis by repressing transposable elements and preventing their mobilization, which is essential for the germline integrity (By similarity). Acts via the piRNA metabolic process, which mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins (By similarity). The PIWIL4-piRNA pathway acts in the nucleus and mediates silencing of active transposons: engages with nascent transposable element transcripts and governs the piRNA-directed DNA methylation and subsequent repression of transposons (By similarity). In contrast to PIWIL1 and PIWIL2, does not show endonuclease activity (By similarity). Directly binds piRNAs, a class of 24 to 30 nucleotide RNAs that are generated by a Dicer-independent mechanism and are primarily derived from transposons and other repeated sequence elements (By similarity). Associates with secondary piRNAs antisense and PIWIL2/MILI is required for such association (By similarity). The piRNA process acts upstream of known mediators of DNA methylation (By similarity). Plays a key role in the piRNA amplification loop, also named ping-pong

amplification cycle, by acting as a 'slicer-incompetent' component that loads cleaved piRNAs from the 'slicer-competent' component PIWIL2 and target them on genomic transposon loci in the nucleus (By similarity). May be involved in the chromatin-modifying pathway by inducing 'Lys-9' methylation of histone H3 at some loci (PubMed:[17544373](#)). In addition to its role in germline, PIWIL4 also plays a role in the regulation of somatic cells activities (By similarity). Plays a role in pancreatic beta cell function and insulin secretion (By similarity). Involved in maintaining cell morphology and functional integrity of retinal epithelial through Akt/GSK3alpha/beta signaling pathway (PubMed:[28025795](#)). When overexpressed, acts as an oncogene by inhibition of apoptosis and promotion of cells proliferation in tumors (PubMed:[22483988](#)).


Cellular Location

Nucleus. Cytoplasm Note=Probable component of the meiotic nuage, also named P granule, a germ-cell-specific organelle required to repress transposon activity during meiosis. PIWIL2/MILI is required for nuclear localization (By similarity). {ECO:0000250|UniProtKB:Q8CGT6}


Tissue Location

Ubiquitously expressed (PubMed:17544373, PubMed:22483988, PubMed:25038252, PubMed:28025795, PubMed:28711973) Detected in retina, retinal pigment epithelia cells (RPE) (at protein level) (PubMed:28025795).

Images




Western blot analysis of HIWI2 expression in PC3 (A), PANC1 (B) whole cell lysates. (Predicted band size: 97 kD; Observed band size: 110 kD)



Immunohistochemical analysis of HIWI2 staining in human rectum cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Flow cytometric analysis of HeLa cells using Anti-HIWI2 Antibody (green) and negative control (red).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.