

Anti-Perforin 1 Antibody

Catalog # AP96015

Product Information

Application	WB, IHC, FC
Primary Accession	P14222
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Calculated MW	61377

Additional Information


Gene ID	5551
Other Names	PFP; Perforin-1; P1; Cytolysin; Lymphocyte pore-forming protein; PFP
Target/Specificity	Recombinant fusion protein of human Perforin 1 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200)
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	PRF1
Synonyms	PFP
Function	Pore-forming protein that plays a key role in granzyme- mediated programmed cell death, and in defense against virus-infected or neoplastic cells (PubMed:20889983, PubMed:21037563, PubMed:24558045, PubMed:9058810, PubMed:9164947). Plays an important role in killing other cells that are recognized as non-self by the immune system, e.g. in transplant rejection or some forms of autoimmune disease (PubMed:9058810). Can insert into the membrane of target cells in its calcium-bound form, oligomerize and form large pores (PubMed:20889983, PubMed:21037563). Promotes cytolysis and apoptosis of target cells by mediating the passage and uptake of cytotoxic granzymes (PubMed:20038786, PubMed:20225066, PubMed:24558045, PubMed:32299851). Facilitates the delivery of cationic cargo protein, while anionic or neural proteins are not delivered efficiently (PubMed:24558045). Perforin pores allow the release of mature caspase-7 (CASP7) into the extracellular milieu (By similarity).
Cellular Location	Cytolytic granule. Secreted. Cell membrane; Multi-pass membrane protein.

Endosome lumen. Note=Stored in cytolytic granules of cytolytic T-lymphocytes and secreted into the cleft between T- lymphocyte and target cell (PubMed:20038786). Inserts into the cell membrane of target cells and forms pores (PubMed:20889983). Membrane insertion and pore formation requires a major conformation change (PubMed:20889983). May be taken up via endocytosis involving clathrin- coated vesicles and accumulate in a first time in large early endosomes (PubMed:20038786).


Images



Western blot analysis of Perforin 1 expression in mouse liver (A) whole cell lysates. (Predicted band size: 61 kD; Observed band size: 61 kD)



Immunohistochemical analysis of Perforin 1 staining in human rectum cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Flow cytometric analysis of Jurkat cells using Anti-Perforin 1 Antibody (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.