

Anti-RAD21 Antibody

Catalog # AP96025

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	O60216
Reactivity	Human, Rat, Mk
Host	Mouse
Clonality	Monoclonal
Calculated MW	71690

Additional Information

Gene ID	5885
Other Names	HR21; KIAA0078; NXP1; Double-strand-break repair protein rad21 homolog; hHR21; Nuclear matrix protein 1; NXP-1; SCC1 homolog
Target/Specificity	Recombinant fusion protein of human RAD21 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200) IF/IC~~N/A
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	RAD21
Function	[Double-strand-break repair protein rad21 homolog]: As a member of the cohesin complex, involved in sister chromatid cohesion from the time of DNA replication in S phase to their segregation in mitosis, a function that is essential for proper chromosome segregation, post-replicative DNA repair, and the prevention of inappropriate recombination between repetitive regions (PubMed: 11509732). The cohesin complex may also play a role in spindle pole assembly during mitosis (PubMed: 11590136). In interphase, cohesins may function in the control of gene expression by binding to numerous sites within the genome (By similarity). May control RUNX1 gene expression (Probable). Binds to and represses APOB gene promoter (PubMed: 25575569). May play a role in embryonic gut development, possibly through the regulation of enteric neuron development (By similarity).
Cellular Location	[Double-strand-break repair protein rad21 homolog]: Nucleus. Nucleus matrix Chromosome Chromosome, centromere. Cytoplasm, cytoskeleton, spindle pole. Note=Associates with chromatin (PubMed:11073952, PubMed:11590136). Before prophase, scattered along chromosome arms

(PubMed:11073952). During prophase and prometaphase, most cohesins dissociate from the arms of condensing chromosome, possibly through PLK1-mediated phosphorylation (PubMed:11931760). A small amount of cohesin remains in centromeric regions and is removed from chromosomes only at the onset of anaphase. At anaphase, cleavage by separase/ESPL1 leads to the dissociation of cohesin from chromosomes and chromosome separation (PubMed:11073952, PubMed:11509732)


Tissue Location

Expressed in the gut (at protein level).


Images



Western blot analysis of RAD21 expression in Hela (A), HEK293 (B), K562 (C), C6 (D), *** (E), COS7 (F) whole cell lysates. (Predicted band size: 72 kD; Observed band size: 72 kD)




Immunohistochemical analysis of RAD21 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of RAD21 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Flow cytometric analysis of HeLa cells using Anti-RAD21 Antibody (green) and negative control (red).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.