

Anti-RbAp46 Antibody

Catalog # AP96035

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	Q16576
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Calculated MW	47820


Additional Information

Gene ID	5931
Other Names	RBAP46; Histone-binding protein RBBP7; Histone acetyltransferase type B subunit 2; Nucleosome-remodeling factor subunit RBAP46; Retinoblastoma-binding protein 7; RBBP-7; Retinoblastoma-binding protein p46
Target/Specificity	Recombinant fusion protein of human RbAp46 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200) IF/IC~~N/A
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	RBBP7
Synonyms	RBAP46
Function	Core histone-binding subunit that may target chromatin remodeling factors, histone acetyltransferases and histone deacetylases to their histone substrates in a manner that is regulated by nucleosomal DNA. Component of several complexes which regulate chromatin metabolism. These include the type B histone acetyltransferase (HAT) complex, which is required for chromatin assembly following DNA replication; the core histone deacetylase (HDAC) complex, which promotes histone deacetylation and consequent transcriptional repression; the nucleosome remodeling and histone deacetylase complex (the NuRD complex), which promotes transcriptional repression by histone deacetylation and nucleosome remodeling; and the PRC2/EED-EZH2 complex, which promotes repression of homeotic genes during development; and the NURF (nucleosome remodeling factor) complex.


Images




Western blot analysis of RbAp46 expression in Jurkat (A), HepG2 (B), F9 (C), C6 (D), LNCAP (E), HL60 (F), HeLa (G), SHSY5Y (H) whole cell lysates. (Predicted band size: 48 kD; Observed band size: 45 kD)



Immunohistochemical analysis of RbAp46 staining in human bladder cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of RbAp46 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



Flow cytometric analysis of HeLa cells using Anti-RbAp46 Antibody (green) and negative control (red).