

Anti-Tyrosine Hydroxylase Antibody

Catalog # AP96076

Product Information

Application	WB, IHC, FC, IF/IC
Primary Accession	P07101
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Calculated MW	58600

Additional Information

Gene ID	7054
Other Names	TYH; Tyrosine 3-monooxygenase; Tyrosine 3-hydroxylase; TH
Target/Specificity	Recombinant fusion protein of human Tyrosine Hydroxylase expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) IHC~~1:100~500 FC~~FC (1/100 - 1/200) IF/IC~~N/A
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	TH (HGNC:11782)
Synonyms	TYH
Function	Catalyzes the conversion of L-tyrosine to L- dihydroxyphenylalanine (L-Dopa), the rate-limiting step in the biosynthesis of catecholamines, dopamine, noradrenaline, and adrenaline. Uses tetrahydrobiopterin and molecular oxygen to convert tyrosine to L-Dopa (PubMed: 15287903 , PubMed: 1680128 , PubMed: 17391063 , PubMed: 24753243 , PubMed: 34922205 , PubMed: 8528210 , Ref.18). In addition to tyrosine, is able to catalyze the hydroxylation of phenylalanine and tryptophan with lower specificity (By similarity). Positively regulates the regression of retinal hyaloid vessels during postnatal development (By similarity).
Cellular Location	Cytoplasm, perinuclear region {ECO:0000250 UniProtKB:P24529}. Nucleus {ECO:0000250 UniProtKB:P04177} Cell projection, axon {ECO:0000250 UniProtKB:P24529}. Cytoplasm {ECO:0000250 UniProtKB:P04177}. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250 UniProtKB:P04177}. Note=When

phosphorylated at Ser-19 shows a nuclear distribution and when phosphorylated at Ser-31 as well at Ser-40 shows a cytosolic distribution (By similarity). Expressed in dopaminergic axons and axon terminals. {ECO:0000250|UniProtKB:P04177}


Tissue Location

Mainly expressed in the brain and adrenal glands.


Images



Western blot analysis of Tyrosine Hydroxylase expression in SHSY5Y (A), PC12 (B) whole cell lysates. (Predicted band size: 59 kD; Observed band size: 59 kD)




Immunohistochemical analysis of Tyrosine Hydroxylase staining in human cervical cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Tyrosine Hydroxylase staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Flow cytometric analysis of HeLa cells using Anti-Tyrosine Hydroxylase Antibody (green) and negative control (red).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.