

Anti-TRAF2 Antibody

Catalog # AP96089

Product Information

Application	WB, FC
Primary Accession	Q12933
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	55859

Additional Information

Gene ID	7186
Other Names	TRAP3; TNF receptor-associated factor 2; E3 ubiquitin-protein ligase TRAF2; Tumor necrosis factor type 2 receptor-associated protein 3
Target/Specificity	Recombinant fusion protein of human TRAF2 expressed in E. Coli
Dilution	WB~~WB (1/500 - 1/1000) FC~~FC (1/100 - 1/200)
Format	Mouse IgG2b. Liquid in PBS, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	TRAF2 {ECO:0000303 PubMed:28489822, ECO:0000312 HGNC:HGNC:12032}
Function	E3 ubiquitin-protein ligase that regulates activation of NF- kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed:10346818, PubMed:11784851, PubMed:12917689, PubMed:15383523, PubMed:18981220, PubMed:19150425, PubMed:19810754, PubMed:19918265, PubMed:19937093, PubMed:20047764, PubMed:20064526, PubMed:20385093, PubMed:20577214, PubMed:22212761). Catalyzes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, IKBKE, MLST8, RIPK1 and TICAM1 (PubMed:23453969, PubMed:28489822). Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases (PubMed:15383523, PubMed:18981220). Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain (PubMed:11907583, PubMed:19506082). Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR (PubMed:15121867). In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE

(PubMed:[23453969](#)). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked ubiquitination of MLST8, thereby inhibiting formation of the mTORC2 complex, while facilitating assembly of the mTORC1 complex (PubMed:[28489822](#)). Required for normal antibody isotype switching from IgM to IgG (By similarity).


Cellular Location

Cytoplasm

Images



Western blot analysis of TRAF2 expression in MCF7 (A), A431 (B), HeLa (C), Jurkat (D), HEK293 (E), Ramos (F) whole cell lysates. (Predicted band size: 56 kD; Observed band size: 55 kD)



Flow cytometric analysis of HeLa cells using Anti-TRAF2 Antibody (green) and negative control (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.