

Anti-GSTP1 Antibody

Catalog # AP96132

Product Information


Application	WB, IHC, FC, IP, IF/IC
Primary Accession	P09211
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23356

Additional Information


Gene ID	2950
Other Names	FAES3; GST3; Glutathione S-transferase P; GST class-pi; GSTP1-1
Target/Specificity	Recombinant fusion protein of human GSTP1
Dilution	WB~~WB (1/500 - 1/2000) IHC~~1:100~500 FC~~1:10~50 IP~~N/A IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	GSTP1 (HGNC:4638)
Synonyms	FAES3, GST3
Function	Catalyzes conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles (PubMed: 1540159 , PubMed: 1567427 , PubMed: 8433974). Involved in the formation of glutathione conjugates of both prostaglandin A2 (PGA2) and prostaglandin J2 (PGJ2) (PubMed: 9084911). Participates in the formation of novel hepxilin regioisomers (PubMed: 21046276). Acts as a negative regulator of ferroptosis by mediating glutathione conjugation and detoxification of 4-hydroxynonenal (4-HNE) reactive aldehyde (PubMed: 38016474). Negatively regulates CDK5 activity via p25/p35 translocation to prevent neurodegeneration (PubMed: 21668448).
Cellular Location	Cytoplasm. Mitochondrion. Nucleus. Note=The 83 N-terminal amino acids function as un uncleaved transit peptide, and arginine residues within it are crucial for mitochondrial localization



Western blot analysis of GSTP1 expression in HEK293T (A), HepG2 (B), mouse liver (C), mouse lung (D) whole cell lysates. (Predicted band size: 23 kD; Observed band size: 25 kD)



Immunohistochemical analysis of GSTP1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of GSTP1 staining in L929 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.