

FOXA3 Rabbit pAb

Catalog # AP96222

Product Information

Application	WB
Primary Accession	P55318
Other Accession	P55318
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37140
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human FOXA3
Epitope Specificity	51-150/350
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 fork-head DNA-binding domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the forkhead class of DNA-binding proteins. These hepatocyte nuclear factors are transcriptional activators for liver-specific transcripts such as albumin and transthyretin, and they also interact with chromatin. Similar family members in mice have roles in the regulation of metabolism and in the differentiation of the pancreas and liver. The crystal structure of a similar protein in rat has been resolved. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	3171
Other Names	FKHH3; HNF3G; TCF3G; HNF3-G; Hnf-3g; Tcf-3g; FOXA3_BOVIN; FOXA3; HNF-3-gamma; Forkhead box protein A3; FOXA3_HUMAN; Fork head-related protein FKH H3; Transcription factor 3G (TCF-3G); FOXA3_MOUSE; FOXA3_RAT; forkhead box A3; hepatocyte nuclear factor 3, gamma
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	FOXA3
-------------	-------

Synonyms	HNH3G, TCF3G
Function	Transcription factor that is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites (By similarity). Originally described as a transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis; binds to and activates transcription from the G6PC1 promoter. Binds to the CYP3A4 promoter and activates its transcription in cooperation with CEBPA. Binds to the CYP3A7 promoter together with members of the CTF/NF-I family. Involved in regulation of neuronal-specific transcription. May be involved in regulation of spermatogenesis.
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00089, ECO:0000269 PubMed:19706729}
Tissue Location	Expressed in erythroleukemia and hepatoma cell lines and in liver and pancreas. Not expressed in any other cell lines or tissues examined.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.