

HEPACAM2 Rabbit pAb

Catalog # AP96271

Product Information

Application	WB, E
Primary Accession	A8MVW5
Other Accession	A8MVW5
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51407
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HEPACAM2
Epitope Specificity	201-300/462
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 2 Ig-like C2-type (immunoglobulin-like) domains.
Post-translational modifications	Poly-ADP-ribosylated (PARsylated) by tankyrase TNKS during late G2 and prophase, leading to translocation to mitotic centrosomes. N-glycosylated.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Widely expressed.

Additional Information

Gene ID	253012
Other Names	MIKI; RGD1305697; HECA2_HUMAN; HEPACAM2; Mitotic kinetics regulator; HECA2_MOUSE; HEPACAM family member 2
Dilution	WB=1:500-2000,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	HEPACAM2
Synonyms	MIKI
Function	Required during prometaphase for centrosome maturation. Following poly-ADP-ribosylation (PARsylation) by TNKS, translocates from the Golgi apparatus to mitotic centrosomes and plays a key role in the formation of

robust microtubules for prompt movement of chromosomes: anchors AKAP9/CG-NAP, a scaffold protein of the gamma- tubulin ring complex and promotes centrosome maturation.

Cellular Location

Golgi apparatus membrane; Single-pass type I membrane protein. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Midbody. Note=In interphase, localizes to the Golgi apparatus. Localizes to centrosomes and spindles during prophase, prometaphase, and metaphase of mitosis, and to midbodies at telophase Translocation to mitotic centrosomes is the result of poly-ADP- ribosylation (PARsylation).

Tissue Location

Widely expressed..

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.