

ANGPTL8 Mouse mAb

Catalog # AP96284

Product Information

Application	WB
Host	Mouse
Clonality	Monoclonal
Calculated MW	20 KDa
Physical State	Liquid
Immunogen	Recombinant mouse ANGPTL8
Epitope Specificity	16-198/198
Isotype	IgG
Purity	affinity purified by Protein G

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SIMILARITY Belongs to the ANGPTL8 family.

SUBUNIT Interacts with ANGPTL3.

Post-translational modifications Proteolytically cleaved at the N-terminus.

DISEASE

Diabetes mellitus, insulin-dependent (IDDM) [MIM:222100]: A multifactorial disorder of glucose homeostasis that is characterized by susceptibility to ketoacidosis in the absence of insulin therapy. Clinical features are polydipsia, polyphagia and polyuria which result from hyperglycemia-induced osmotic diuresis and secondary thirst. These derangements result in long-term complications that affect the eyes, kidneys, nerves, and blood vessels. {ECO:0000305 | PubMed:24078058}. Note=The gene represented in this entry may be involved in disease pathogenesis. Increased protein levels are observed in the serum of patients. This result should however be reinvestigated in light of recent advances that suggest that this protein is not promoting pancreatic beta cell proliferation. Diabetes mellitus, non-insulin-dependent (NIDDM) [MIM:125853]: A multifactorial disorder of glucose homeostasis caused by a lack of sensitivity to the body's own insulin. Affected individuals usually have an obese body habitus and manifestations of a metabolic syndrome characterized by diabetes, insulin resistance, hypertension and hypertriglyceridemia. The disease results in long-term complications that affect the eyes, kidneys, nerves, and blood vessels. {ECO:0000305 | PubMed:24852694, ECO:0000305 | PubMed:24963292, ECO:0000305 | PubMed:25024395, ECO:0000305 | PubMed:25303484}. Note=The gene represented in this entry may be involved in disease pathogenesis. Increased protein levels are observed in the serum of patients and are associated with insulin resistance (PubMed:25024395, PubMed:25303484, PubMed:24963292, PubMed:24852694). According to another report, protein levels are decreased in the serum of patients (PubMed:25050901). Discrepancies between increased and decreased levels of proteins levels in NIDDM patients may be explained by the use of different kits developed on the market that either use antibodies recognizing the N-terminal or the C-terminal part of the protein (PubMed:25099942). These results should however be reinvestigated in light of recent advances that suggest that this protein is not promoting pancreatic beta cell proliferation. This product as supplied is intended for research use only, not for use in

Important Note

Background Descriptions	human, therapeutic or diagnostic applications. ANGPTL8 (Angiopoietin Like 8) is a Protein Coding gene. Diseases associated with ANGPTL8 include Breast Angiosarcoma and Breast Sarcoma. Among its related pathways are Lipoprotein metabolism and Metabolism.
Additional Information	
Other Names	C19orf80; PRO1185; PVPA599; RIFL; TD26; ANGL8_HUMAN; ANGPTL8; Betatrophin; Lipasin; Refeeding-induced fat and liver protein; angiopoietin like 8; chromosome 19 open reading frame 80; angiopoietin-like 8
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Name	ANGPTL8 Mouse mAb
Function	Hormone that acts as a blood lipid regulator by regulating serum triglyceride levels (PubMed:22569073, PubMed:22809513, PubMed:23150577). May be involved in the metabolic transition between fasting and refeeding; required to direct fatty acids to adipose tissue for storage in the fed state.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.