

KRT72 Rabbit pAb

Catalog # AP96357

Product Information

Application	WB
Primary Accession	Q14CN4
Other Accession	Q14CN4
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55877
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KRT72
Epitope Specificity	431-511/511
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the intermediate filament family.
SUBUNIT	Heterotetramer of two type I and two type II keratins.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells. The type II keratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This gene encodes a type II keratin that is specifically expressed in the inner root sheath of hair follicles. The type II keratins are clustered in a region of chromosome 12q12-q13. Multiple transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jun 2009]

Additional Information

Gene ID	140807
Other Names	CK-72; K6IRS2; K6irs; K72; KRT6; KRT6IRS2; K2C72_HUMAN; KRT72; Cytokeratin-72 (CK-72); Keratin-72 (K72); Type II inner root sheath-specific keratin-K6irs2; Type-II keratin Kb35; KB35; keratin 72; keratin 72, type II
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	KRT72
Synonyms	K6IRS2, KB35, KRT6, KRT6IRS2
Function	Has a role in hair formation. Specific component of keratin intermediate filaments in the inner root sheath (IRS) of the hair follicle (Probable).
Tissue Location	Highly expressed in hair follicles from scalp and eyebrow. Also expressed in palmoplantar epidermis. Not expressed in face skin despite the presence of fine hairs histologically. In hair, it is specifically present in the inner root sheath (IRS) of the hair follicle. Present in the IRS cuticle, but not in Henle or Huxley layers of the IRS. In the IRS cuticle, its presence is delayed up to the height of the apex of the dermal papilla (at protein level)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.