

TBXT Recombinant Mouse mAb

Catalog # AP96362

Product Information

Application	WB
Host	Mouse
Clonality	Recombinant
Calculated MW	48 KDa
Physical State	Liquid
Isotype	IgG1, Kappa
Purity	affinity purified by Protein G
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 T-box DNA-binding domain.
SUBUNIT	Monomer.
DISEASE	Neural tube defects (NTD) [MIM:182940]: Congenital malformations of the central nervous system and adjacent structures related to defective neural tube closure during the first trimester of pregnancy. Failure of neural tube closure can occur at any level of the embryonic axis. Common NTD forms include anencephaly, myelomeningocele and spina bifida, which result from the failure of fusion in the cranial and spinal region of the neural tube. NTDs have a multifactorial etiology encompassing both genetic and environmental components. Note=Disease susceptibility is associated with variations affecting the gene represented in this entry. Chordoma (CHDM) [MIM:215400]: Rare, clinically malignant tumors derived from notochordal remnants. They occur along the length of the spinal axis, predominantly in the sphenooccipital, vertebral and sacrococcygeal regions. They are characterized by slow growth, local destruction of bone, extension into adjacent soft tissues and rarely, distant metastatic spread. Note=Disease susceptibility is associated with variations affecting the gene represented in this entry. Susceptibility to development of chordomas is due to a T gene duplication.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is an embryonic nuclear transcription factor that binds to a specific DNA element, the palindromic T-site. It binds through a region in its N-terminus, called the T-box, and effects transcription of genes required for mesoderm formation and differentiation. The protein is localized to notochord-derived cells. Variation in this gene was associated with susceptibility to neural tube defects and chordoma. A mutation in this gene was found in a family with sacral agenesis with vertebral anomalies. [provided by RefSeq, Sep 2018]

Additional Information

Other Names	SAVA; T; TFT; Bra; D17Mit170; Low; Lr; T1; Tbxt; TI2; TI3; cou; me75; TBXT_HUMAN; Brachyury protein; Protein T; TBXT_MOUSE; T-box transcription factor T; T, brachyury homolog (mouse); T brachyury transcription factor
--------------------	--

Dilution	WB=1:500-2500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Name	TBXT Recombinant Mouse mAb
Function	Involved in the transcriptional regulation of genes required for mesoderm formation and differentiation. Binds to a palindromic site (called T site) and activates gene transcription when bound to such a site.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.