10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# phospho-RPS6KA1 (Thr359) Rabbit pAb

Catalog # AP96395

## Product Information

| | |
|---|---|
| **Application** | WB, E |
| **Reactivity** | Human |
| **Predicted** | Rat, Pig, Mouse, Rabbit, Chicken, Dog, Horse, Sheep |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Calculated MW** | 81 KDa |
| **Physical State** | Liquid |
| **Immunogen** | KLH conjugated Synthesised phosphopeptide derived from human RPS6KA1 around the phosphorylation site of Thr359 |
| **Epitope Specificity** | TS(P-S)PI |
| **Isotype** | IgG |
| **Purity** | affinity purified by Protein A |
| **Buffer** | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| **SIMILARITY** | Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily.Contains 1 AGC-kinase C-terminal domain.Contains 2 protein kinase domains. |
| **SUBUNIT** | Forms a complex with either MAPK1/ERK2 or MAPK3/ERK1 in quiescent cells. Transiently dissociates following mitogenic stimulation. Interacts with ETV1/ER81 and FGFR1. |
| **Post-translational modifications** | Activated by phosphorylation at Ser-221 by PDPK1. Autophosphorylated on Ser-380, as part of the activation process. May be phosphorylated at Thr-359 and Ser-363 by MAPK1/ERK2 and MAPK3/ERK1.N-terminal myristoylation results in an activated kinase in the absence of added growth factors. |
| **Important Note** | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |
| **Background Descriptions** | Rsk1 is a member of a family of 90kDa ribosomal protein S6 kinases, which includes Rsk1, Rsk2 and Rsk3. These are broadly expressed serine/threonine protein kinases activated in response to mitogenic stimuli, including extracellular signal regulated protein kinases Erk1 and Erk2. Rsk1 is activated by MAPK in vitro and in vivo via phosphorylation. Active Rsks appear to play a major role in transcriptional regulation by translocating to the nucleus and phosphorylating c Fos and CREB. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 6195 |
| **Other Names** | KS6A1_MOUSE; Rps6ka1; S6K-alpha-1; 90 kDa ribosomal protein S6 kinase 1 (p90-RSK 1 \| p90RSK1 \| p90S6K); MAP kinase-activated protein kinase 1a (MAPK-activated protein kinase 1a \| MAPKAP kinase 1a \| MAPKAPK-1a); Ribosomal S6 kinase 1 (RSK-1); 2.7.11.1; Mapkapk1a; Rsk1; KS6A1_RAT; HU-1; MAPKAPK1; RSK; p90Rsk; ribosomal protein S6 kinase A1; ribosomal protein S6 kinase, 90kD, polypeptide 1; RPS6KA1 \| p90-RSK1 (phospho-T359); |

| | |
|---|---|
| | p-p90-RSK1; phospho-p90-RSK1; p-RPS6KA1; RPS6KA1 \| p90-RSK1 (phospho-Thr359) |
| **Dilution** | WB=1:500-2000,ELISA=1:5000-10000 |
| **Storage** | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C. |
| **Name** | phospho-RPS6KA1 (Thr359) Rabbit pAb |
| **Function** | Serine/threonine-protein kinase that acts downstream of ERK (MAPK1/ERK2 and MAPK3/ERK1) signaling and mediates mitogenic and stress-induced activation of the transcription factors CREB1, ETV1/ER81 and NR4A1/NUR77, regulates translation through RPS6 and EIF4B phosphorylation, and mediates cellular proliferation, survival, and differentiation by modulating mTOR signaling and repressing pro-apoptotic function of BAD and DAPK1. In fibroblast, is required for EGF-stimulated phosphorylation of CREB1, which results in the subsequent transcriptional activation of several immediate-early genes. In response to mitogenic stimulation (EGF and PMA), phosphorylates and activates NR4A1/NUR77 and ETV1/ER81 transcription factors and the cofactor CREBBP. Upon insulin-derived signal, acts indirectly on the transcription regulation of several genes by phosphorylating GSK3B at 'Ser-9' and inhibiting its activity. Phosphorylates RPS6 in response to serum or EGF via an mTOR-independent mechanism and promotes translation initiation by facilitating assembly of the preinitiation complex. In response to insulin, phosphorylates EIF4B, enhancing EIF4B affinity for the EIF3 complex and stimulating cap-dependent translation. Is involved in the mTOR nutrient-sensing pathway by directly phosphorylating TSC2 at 'Ser-1798', which potently inhibits TSC2 ability to suppress mTOR signaling, and mediates phosphorylation of RPTOR, which regulates mTORC1 activity and may promote rapamycin-sensitive signaling independently of the PI3K/AKT pathway. Mediates cell survival by phosphorylating the pro-apoptotic proteins BAD and DAPK1 and suppressing their pro-apoptotic function. Promotes the survival of hepatic stellate cells by phosphorylating CEBPB in response to the hepatotoxin carbon tetrachloride (CCl4). Is involved in cell cycle regulation by phosphorylating the CDK inhibitor CDKN1B, which promotes CDKN1B association with 14-3-3 proteins and prevents its translocation to the nucleus and inhibition of G1 progression. |

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.