10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

WISP2 Rabbit pAb

Catalog # AP96431

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	O76076
Other Accession	O76076
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26825
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human WISP2
Epitope Specificity	101-200/250
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the CCN family. Contains 1 IGFBP N-terminal domain. Contains 1 TSP type-1 domain. Contains 1 VWFC domain.
SUBUNIT	Secreted
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	WISP2 a member of the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like (CT) domain. WISP2 lacks the CT domain which is implicated in dimerization and heparin binding. It is 72% identical to the mouse protein at the amino acid level. This gene may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. Its expression in colon tumors is reduced while the other two WISP members are overexpressed in colon tumors. It is expressed at high levels in bone tissue, and may play an important role in modulating bone turnover.

Additional Information

Gene ID	8839
Other Names	CT58; CTGF-L; WISP2; CCN5_HUMAN; CCN5; Connective tissue growth factor-like protein (CTGF-L); Connective tissue growth factor-related protein 58; WNT1-inducible-signaling pathway protein 2 (WISP-2); CTGFL; cellular communication network factor 5; WNT1 inducible signaling pathway protein 2
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-

	500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CCN5 (HGNC:12770)
Synonyms	CT58, CTGFL, WISP2
Function	May play an important role in modulating bone turnover. Promotes the adhesion of osteoblast cells and inhibits the binding of fibrinogen to integrin receptors. In addition, inhibits osteocalcin production.
Cellular Location	Secreted.
Tissue Location	Expressed in primary osteoblasts, fibroblasts, ovary, testes, and heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.