10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

C9orf95 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9650a

Product Information

Application	WB, E
Primary Accession	Q9NWW6
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB24277
Calculated MW	23193
Antigen Region	24-50

Additional Information

Gene ID	54981
Other Names	Nicotinamide riboside kinase 1, NRK 1, NmR-K 1, Nicotinic acid riboside kinase 1, Ribosylnicotinamide kinase 1, RNK 1, Ribosylnicotinic acid kinase 1, NMRK1, C9orf95, NRK1
Target/Specificity	This C9orf95 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 24-50 amino acids from the N-terminal region of human C9orf95.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	C9orf95 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NMRK1
Synonyms	C9orf95, NRK1

Function Catalyzes the phosphorylation of nicotinamide riboside (NR) and nicotinic acid riboside (NaR) to form nicotinamide mononucleotide (NMN) and nicotinic acid mononucleotide (NaMN). The enzyme also phosphorylates the antitumor drugs tiazofurin and 3-deazaguanosine.


Background

Nicotinamide adenine dinucleotide (NAD+) is essential for life in all organisms, both as a coenzyme for oxidoreductases and as a source of ADP-ribosyl groups used in various reactions. Nicotinic acid and nicotinamide, collectively known as niacin, are the vitamin precursors of NAD+. Nicotinamide riboside kinases, such as NRK1, function to synthesize NAD+ through nicotinamide mononucleotide using nicotinamide riboside as the precursor (Bieganowski and Brenner, 2004 [PubMed 15137942]).[supplied by OMIM].


References

Khan, J.A., et al. Structure 15(8):1005-1013(2007)
Humphray, S.J., et al. Nature 429(6990):369-374(2004)
Bieganowski, P., et al. Cell 117(4):495-502(2004)

Images



All lanes : Anti-C9orf95 Antibody (N-term) at 1:1000 dilution Lane 1: WiDr whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size : 23kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-C9orf95 Antibody (N-term) at 1:1000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: RPMI 8226 whole cell lysate Lane 3: M. kidney whole cell lysate Lane 4: R. brain whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size : 23kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.