

AMACR Rabbit pAb

Catalog # AP96517

Product Information

Application	WB
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42 KDa
Physical State	Liquid
Immunogen	Recombinant Mouse AMACR protein
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the CaiB/BaiF CoA-transferase family.
DISEASE	Alpha-methylacyl-CoA racemase deficiency (AMACRD) [MIM:614307]: A rare autosomal recessive peroxisomal disorder characterized by elevated plasma concentrations of pristanic acid C27-bile-acid intermediates, and adult onset of variable neurodegenerative symptoms affecting the central and peripheral nervous systems. Features may include seizures, visual failure, sensorimotor neuropathy, spasticity, migraine, and white matter hyperintensities on brain imaging. Note=The disease is caused by mutations affecting the gene represented in this entry. Congenital bile acid synthesis defect 4 (CBAS4) [MIM:214950]: A disorder characterized by the presence of trihydroxycoprostanic acid in the bile and absence of cholic acid. Patients manifest neonatal jaundice, intrahepatic cholestasis and bile duct deficiency. Note=The disease is caused by mutations affecting the gene represented in this entry.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy, pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the upstream neighboring C1QTNF3 (C1q and tumor necrosis factor related protein 3) gene. [provided by RefSeq, Mar 2011]

Additional Information

Other Names	AMACRD; CBAS4; P504S; RACE; RM; AMACR_HUMAN; AMACR; 2-methylacyl-CoA racemase; 5.1.99.4; alpha-methylacyl-CoA racemase
Dilution	WB=1:500-2000

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Name	AMACR Rabbit pAb
Function	Racemization of 2-methyl-branched fatty acid CoA esters. Responsible for the conversion of pristanoyl-CoA and C27-bile acyl-CoAs to their (S)-stereoisomers.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.