

phospho-PKA beta (Ser339) Rabbit pAb

Catalog # AP96546

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40 kDa
Physical State	Liquid
Immunogen	KLH conjugated Synthesised phosphopeptide derived from human PKA beta around the phosphorylation site of Ser339
Epitope Specificity	RV(p-S)IT
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. cAMP subfamily. Contains 1 AGC-kinase C-terminal domain. Contains 1 protein kinase domain.
SUBUNIT	Belongs to the cAMP-dependent kinase regulatory chain family. Contains 2 cyclic nucleotide-binding domains.
Post-translational modifications	Asn-3 is partially deaminated to Asp giving rise to 2 major isoelectric variants, called CB and CA respectively (By similarity).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-dependent protein kinase, which transduces the signal through phosphorylation of different target proteins. The inactive kinase holoenzyme is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. The protein encoded by this gene is a member of the Ser/Thr protein kinase family and is a catalytic subunit of cAMP-dependent protein kinase. Several alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jun 2011].

Additional Information

Gene ID	5567
Other Names	KAPCB_HUMAN; PRKACB; PKA C-beta; 2.7.11.11; CAFD2; PKACB; protein kinase cAMP-activated catalytic subunit beta; protein kinase, cAMP-dependent, catalytic, beta; protein kinase, cAMP-dependent, beta catalytic subunit; cAMP-dependent protein kinase catalytic subunit beta; PRKACB PKA beta (phospho-S339); p-PKA beta; phospho-PKA beta; p-PKACB; PRKACB PKA beta (phospho-Ser339)

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Name	phospho-PKA beta (Ser339) Rabbit pAb
Function	Four types of regulatory chains are found: I-alpha, I-beta, II-alpha, and II-beta. Their expression varies among tissues and is in some cases constitutive and in others inducible.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.