10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TEAD1 Recombinant Rabbit mAb

Catalog # AP96676

Product Information

Application	WB
Primary Accession	P28347, Q15561, Q15562, Q99594
Other Accession	P28347, Q15561, Q15562, Q99594
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Recombinant
Calculated MW	47946
Physical State	Liquid
Immunogen	A synthesized peptide derived from human TEAD1
Epitope Specificity	1-45
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol.
SUBUNIT	Interacts with YAP1 and WWTR1/TAZ.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Transcription factor which plays a key role in the Hippo signaling pathway, a pathway involved in organ size control and tumor suppression by restricting proliferation and promoting apoptosis.

Additional Information

Gene ID	7003
Other Names	AA; NTEF-1; REF1; TCF-13; TCF13; TEAD-1; TEF-1; TEA domain transcription factor 1; atrophia areata, peripapillary chorioretinal degeneration; TEA domain family member 1 (SV40 transcriptional enhancer factor); transcriptional enhancer factor 1
Dilution	WB=1:1000-2000,ICC/IF=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TEAD1
Synonyms	TCF13, TEF1

Function	Transcription factor which plays a key role in the Hippo signaling pathway, a pathway involved in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein MST1/MST2, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Acts by mediating gene expression of YAP1 and WWTR1/TAZ, thereby regulating cell proliferation, migration and epithelial mesenchymal transition (EMT) induction. Binds specifically and cooperatively to the SPH and GT-IIC 'enhansons' (5'-GTGGAATGT-3') and activates transcription in vivo in a cell-specific manner. The activation function appears to be mediated by a limiting cell-specific transcriptional intermediary factor (TIF). Involved in cardiac development. Binds to the M-CAT motif.
Cellular Location	Nucleus.
Tissue Location	Preferentially expressed in skeletal muscle. Lower levels in pancreas, placenta, and heart

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.