

phospho-ASC (Tyr144) Rabbit pAb

Catalog # AP96692

Product Information

Application	WB
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21 kDa
Physical State	Liquid
Immunogen	Asc (Tyr144) phospho-peptide (coupled to KLH) corresponding to amino acid residues surrounding Tyr-144 in mouse Asc. This peptide sequence is highly conserved in human and rat Asc.
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	PBS + 1 mg/ml BSA, 0.05% NaN3 and 50% glycerol
SIMILARITY	Contains 1 CARD domain. Contains 1 DAPIN domain.
SUBUNIT	Forms complexes with other DAPIN domain-containing proteins. Interacts with CIAS1/PYPAF1 and PYDC1.
Post-translational modifications	Phosphorylated.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Promotes caspase-mediated apoptosis. This proapoptotic activity is mediated predominantly through the activation of caspase-9. May be a component of the inflammasome, a protein complex which also includes NALP2, CARD8 and CASP1 and whose function would be the activation of proinflammatory caspases. Tissue specificity: Widely expressed at low levels. Detected in peripheral blood leukocytes, lung, small intestine, spleen, thymus, colon and at lower levels in placenta, liver and kidney. Very low expression in skeletal muscle, heart and brain. Detected in the leukemia cell lines HL-60 and U937, but not in Jurkat T-cell lymphoma and Daudi Burkitt's lymphoma.

Additional Information

Other Names	ASC; CARD5; TMS; TMS-1; TMS1; ASC_HUMAN; PYCARD; hASC; Caspase recruitment domain-containing protein 5; PYD and CARD domain-containing protein; Target of methylation-induced silencing 1; PYD and CARD domain containing; apoptosis-associated speck-like; apoptosis-associated speck-like protein containing a CARD; Caspase, PYCARD, AIM, ASC, CARD5, TMS1
Dilution	WB=1:300-5000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Name	phospho-ASC (Tyr144) Rabbit pAb

Function

Promotes caspase-mediated apoptosis. This proapoptotic activity is mediated predominantly through the activation of caspase-9. May be a component of the inflammasome, a protein complex which also includes NALP2, CARD8 and CASP1 and whose function would be the activation of proinflammatory caspases.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.