

SUCLG2 Rabbit pAb

Catalog # AP96693

Product Information

Application	WB
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46 kDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SUCLG2
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Background Descriptions	This gene encodes a GTP-specific beta subunit of succinyl-CoA synthetase. Succinyl-CoA synthetase catalyzes the reversible reaction involving the formation of succinyl-CoA and succinate. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 5 and 12. [provided by RefSeq, Apr 2010]

Additional Information

Other Names	succinate-CoA ligase GDP-forming subunit beta; SUCB2_HUMAN; Succinate--CoA ligase [GDP-forming] subunit beta, mitochondrial; GTP-specific succinyl-CoA synthetase subunit beta; G-SCS; GTPSCS; Itaconyl--CoA ligase [GDP-forming] subunit beta; Succinyl-CoA synthetase beta-G chain; Malyl--CoA ligase [GDP-forming] subunit beta; SCS-betaG
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
Name	SUCLG2 Rabbit pAb
Function	GTP-specific succinyl-CoA synthetase functions in the citric acid cycle (TCA), coupling the hydrolysis of succinyl-CoA to the synthesis of GTP and thus represents the only step of substrate-level phosphorylation in the TCA (PubMed:40108300).The beta subunit provides nucleotide specificity of the enzyme and binds the substrate succinate, while the binding sites for coenzyme A and phosphate are found in the alpha subunit (By similarity).Also able to act as a GTP-specific itaconyl- and malyl-CoA synthetase (PubMed:40108300).