

Calmodulin 3 Recombinant Rabbit mAb

Catalog # AP96695

Product Information

Application	WB, IHC-P, IHC-F, IF, IP, E
Primary Accession	P0DP25
Other Accession	P0DP25
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Recombinant
Calculated MW	16838
Isotype	IgG
Purity	affinity purified by Protein A

Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Background Descriptions	This gene encodes a member of a family of proteins that binds calcium and functions as a enzymatic co-factor. Activity of this protein is important in the regulation of the cell cycle and cytokinesis. Multiple alternatively spliced transcript variants have been observed at this gene. [provided by RefSeq, Aug 2016]

Additional Information

Gene ID	801;805;808
Other Names	Calmodulin-3;CALM3;CALML2;CAM3;CAMC;CAMIII;Calmodulin;CaM
Dilution	IHC-P=1:50-200,IHC-F=1:50-200,WB=1:500-2000,IF=1:50-200,ELISA=1:5000-10 000,IP=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CALM3 (HGNC:1449)
Function	Calmodulin acts as part of a calcium signal transduction pathway by mediating the control of a large number of enzymes, ion channels, aquaporins and other proteins through calcium-binding (PubMed: 16760425 , PubMed: 31454269). Calcium-binding is required for the activation of calmodulin (PubMed: 16760425 , PubMed: 31454269 , PubMed: 35568036). Among the enzymes to be stimulated by the calmodulin- calcium complex are a number of protein kinases, such as myosin light- chain kinases and calmodulin-dependent protein kinase type II (CaMK2), and phosphatases

(PubMed:[16760425](#), PubMed:[35568036](#)). Together with CCP110 and centrin, is involved in a genetic pathway that regulates the centrosome cycle and progression through cytokinesis (PubMed:[16760425](#)).

Cellular Location

Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome
Note=Distributed throughout the cell during interphase, but during mitosis becomes dramatically localized to the spindle poles and the spindle microtubules

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.