

HATH6/Protein atonal homolog 8 Rabbit pAb

Catalog # AP96708

Product Information

Application	WB
Primary Accession	Q99NA2
Other Accession	Q99NA2
Reactivity	Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34785
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from mouse HATH6/Protein atonal homolog 8
Epitope Specificity	1-100/322
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 basic helix-loop-helix (bHLH) domain.
SUBUNIT	Efficient DNA binding requires dimerization with another bHLH protein
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Putative transcription factor. May be implicated in specification and differentiation of neuronal cell lineages in the brain. May participate in kidney development and may be involved in podocyte differentiation.

Additional Information

Gene ID	71093
Other Names	HATH6; bHLHa21; 4933425C05Rik; Math6; okadin; RGD1561512; ATOH8_HUMAN; ATOH8; Class A basic helix-loop-helix protein 21 (bHLHa21); Helix-loop-helix protein hATH-6 (hATH6); Protein atonal homolog 8; ATH6; ATOH8_MOUSE; Helix-loop-helix protein mATH-6 (mATH6); atonal bHLH transcription factor 8; atonal homolog 8 (Drosophila); atonal homolog bHLH transcription factor 8; basic helix loop helix transcription factor 6
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	Atoh8 {ECO:0000312 MGI:MGI:1918343}
Synonyms	Ath6
Function	Transcription factor that binds a palindromic (canonical) core consensus DNA sequence 5'-CANNTG- 3' known as an E-box element, possibly as a heterodimer with other bHLH proteins (By similarity). Regulates endothelial cell proliferation, migration and tube-like structures formation (By similarity). Modulates endothelial cell differentiation through NOS3 (By similarity). May be implicated in specification and differentiation of neuronal cell lineages in the brain (PubMed:11733035). May participate in kidney development and may be involved in podocyte differentiation (PubMed:16937370). During early embryonic development is involved in tissue-specific differentiation processes that are dependent on class II bHLH factors and namely modulates the differentiation program initiated by the pro-endocrine factor NEUROG3 (PubMed:18560595). During myogenesis, may play a role during the transition of myoblasts from the proliferative phase to the differentiation phase (PubMed:24186058). Positively regulates HAMP transcription in two ways, firstly by acting directly on the HAMP promoter via E-boxes binding and indirectly through increased phosphorylation of SMAD protein complex (By similarity). Repress NEUROG3-dependent gene activation in a gene-specific manner through at least two mechanisms; requires only either the sequestering of a general partner such as TCF3 through heterodimerization, either also requires binding of the bHLH domain to DNA via a basic motif (PubMed:23938248).
Cellular Location	Nucleus. Nucleus speckle {ECO:0000250 UniProtKB:Q96SQ7}. Cytoplasm
Tissue Location	Expressed by subsets of mature neurons (PubMed:11733035). Expressed in kidney (podocytes) (PubMed:16937370) Expression is restricted to the atria, lung mesenchyme, and vascular smooth muscle (PubMed:23836893).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.