

IL17A Rabbit pAb

Catalog # AP96716

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q62386
Other Accession	Q62386
Reactivity	Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17490
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from mouse IL-17
Epitope Specificity	85-150/150
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the IL-17 family.
SUBUNIT	Homodimer.
Post-translational modifications	Found both in glycosylated and nonglycosylated forms.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	<p>This gene is a member of the IL-17 receptor family which includes five members (IL-17RA-E) and the encoded protein is a proinflammatory cytokine produced by activated T cells. IL-17A-mediated downstream pathways induce the production of inflammatory molecules, chemokines, antimicrobial peptides, and remodeling proteins. The encoded protein elicits crucial impacts on host defense, cell trafficking, immune modulation, and tissue repair, with a key role in the induction of innate immune defenses. This cytokine stimulates non-hematopoietic cells and promotes chemokine production thereby attracting myeloid cells to inflammatory sites. This cytokine also regulates the activities of NF-kappaB and mitogen-activated protein kinases and can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). IL-17A plays a pivotal role in various infectious diseases, inflammatory and autoimmune disorders, and cancer. High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. The lung damage induced by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL17A. [provided by RefSeq, Sep 2020]</p>

Additional Information

Gene ID 16171

Other Names	CTLA-8; CTLA8; IL-17; IL-17A; IL17; ILA17; IL17_HUMAN; IL17A; Cytotoxic T-lymphocyte-associated antigen 8 (CTLA-8); IL17_MOUSE; IL17_RAT; interleukin 17A; interleukin 17 (cytotoxic T-lymphocyte-associated serine esterase 8); cytotoxic T-lymphocyte-associated protein 8
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	Il17a
Synonyms	Ctla8, IL17
Function	<p>Effector cytokine of innate and adaptive immune system involved in antimicrobial host defense and maintenance of tissue integrity (PubMed:18025225, PubMed:19144317, PubMed:26431948). Signals via IL17RA-IL17RC heterodimeric receptor complex, triggering homotypic interaction of IL17RA and IL17RC chains with TRAF3IP2 adapter. This leads to downstream TRAF6-mediated activation of NF-kappa-B and MAPkinase pathways ultimately resulting in transcriptional activation of cytokines, chemokines, antimicrobial peptides and matrix metalloproteinases, with potential strong immune inflammation (PubMed:16200068, PubMed:17911633, PubMed:19144317, PubMed:26431948). Plays an important role in connecting T cell-mediated adaptive immunity and acute inflammatory response to destroy extracellular bacteria and fungi. As a signature effector cytokine of T-helper 17 cells (Th17), primarily induces neutrophil activation and recruitment at infection and inflammatory sites (PubMed:18025225). In airway epithelium, mediates neutrophil chemotaxis via induction of CXCL1 and CXCL5 chemokines (PubMed:18025225, PubMed:27923703). In secondary lymphoid organs, contributes to germinal center formation by regulating the chemotactic response of B cells to CXCL12 and CXCL13, enhancing retention of B cells within the germinal centers, B cell somatic hypermutation rate and selection toward plasma cells (PubMed:18157131). Effector cytokine of a subset of gamma-delta T cells that functions as part of an inflammatory circuit downstream IL1B, TLR2 and IL23A-IL12B to promote neutrophil recruitment for efficient bacterial clearance (PubMed:17372004, PubMed:20364087, PubMed:28709803). Effector cytokine of innate immune cells including invariant natural killer cell (iNKT) and group 3 innate lymphoid cells that mediate initial neutrophilic inflammation (PubMed:17470641, PubMed:23255360). Involved in the maintenance of the integrity of epithelial barriers during homeostasis and pathogen infection. Upon acute injury, has a direct role in epithelial barrier formation by regulating OCLN localization and tight junction biogenesis (PubMed:26431948). As part of the mucosal immune response induced by commensal bacteria, enhances host's ability to resist pathogenic bacterial and fungal infections by promoting neutrophil recruitment and antimicrobial peptides release (PubMed:28709803). In synergy with IL17F, mediates the production of antimicrobial beta-defensins DEFB1, DEFB103A, and DEFB104A by mucosal epithelial cells, limiting the entry of microbes through the epithelial barriers (PubMed:19144317). Involved in antiviral host defense through various mechanisms (PubMed:21946434, PubMed:26735852, PubMed:27795421). Enhances immunity against West Nile virus by promoting T cell cytotoxicity (PubMed:27795421). May play a beneficial role in influenza A virus (H5N1) infection by enhancing B cell recruitment and immune</p>

response in the lung (PubMed:[21946434](#)). Contributes to influenza A virus (H1N1) clearance by driving the differentiation of B-1a B cells, providing for production of virus-specific IgM antibodies at first line of host defense (PubMed:[26735852](#)).

Cellular Location

Secreted.

Tissue Location

Expressed by Th17 cell lineage (at protein level). The expression pattern reflects the differentiation state, with IL17A- IL17F heterodimers produced at higher levels than IL17A-IL17A and IL17F-IL17F dimers in fully differentiated Th17 cells (PubMed:16990136, PubMed:18025225). Expressed in innate lymphoid cells (at protein level) (PubMed:23255360, PubMed:28709803). Expressed in gamma-delta T cell subsets (at protein level) (PubMed:17372004, PubMed:20364087, PubMed:26431948, PubMed:28709803). Expressed in iNKT cells (at protein level) (PubMed:17470641).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.