

DEPDC1 Rabbit pAb

Catalog # AP96754

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q5TB30
Other Accession	Q5TB30
Host	Rabbit
Clonality	Polyclonal
Calculated MW	92960
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DEPDC1
Epitope Specificity	606-660/811
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 DEP domain. Contains 1 Rho-GAP domain.
SUBUNIT	Isoform 2 and isoform 5 can form homodimers and heterodimers. Interacts with ZNF224.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	DEPDC1 is a 784 amino acid nuclear protein expressed in testis and up-regulated in bladder cancer cells. Containing a DEP domain and a Rho-GAP domain, DEPDC1 may play an essential role in the growth of bladder cancer cells, and is considered a novel protein target for bladder cancer therapy. Existing as five isoforms produced by alternative splicing events, DEPDC1 is encoded by a gene located on human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma

Additional Information

Gene ID	55635
Other Names	DEP.8; DEPDC1-V2; DEPDC1A; SDP35; DEP1A_HUMAN; DEPDC1; DEP domain containing 1
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	DEPDC1
Synonyms	DEPDC1A
Function	May be involved in transcriptional regulation as a transcriptional corepressor. The DEPDC1A-ZNF224 complex may play a critical role in bladder carcinogenesis by repressing the transcription of the A20 gene, leading to transport of NF-KB protein into the nucleus, resulting in suppression of apoptosis of bladder cancer cells.
Cellular Location	Nucleus. Note=Colocalizes with ZNF224 at the nucleus
Tissue Location	Expressed in testis. Up-regulated in bladder cancer cells (at protein level).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.