10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PP2A alpha + beta Rabbit pAb

Catalog # AP96762

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P62714
Other Accession	P62714
Reactivity	Rat, Human
Predicted	Pig, Mouse, Rabbit, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35575
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human PP2A beta/PPP2CB
Epitope Specificity	21-120/309
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the PPP phosphatase family. PP-1 subfamily.
SUBUNIT	PP2A consists of a common heterodimeric core enzyme, composed of PPP2CA a 36 kDa catalytic subunit (subunit C) and PPP2R1A a 65 kDa constant regulatory subunit (PR65 or subunit A), that associates with a variety of regulatory subunits. Proteins that associate with the core dimer include three families of regulatory subunits B (the R2/B/PR55/B55, R3/B''/PR72/PR130/PR59 and R5/B'/B56 families), the 48 kDa variable regulatory subunit, viral proteins, and cell signaling molecules. Interacts with NXN; the interaction is direct (By similarity). Interacts with TP53, SGOL1 and SGOL2. Interacts with AXIN1; the interaction dephosphorylates AXIN1. Interacts with PIM3; this interaction promotes dephosphorylation, ubiquitination and proteasomal degradation of PIM3. Interacts with RAF1.
Post-translational modifications	Reversibly methyl esterified on Leu-309. Carboxyl methylation may play a role in holoenzyme assembly, enhancing the affinity of the PP2A core enzyme for some, but not all, regulatory subunits. It varies during the cell cycle. Phosphorylation of either threonine (by autophosphorylation-activated protein kinase) or tyrosine results in inactivation of the phosphatase. Auto-dephosphorylation has been suggested as a mechanism for reactivation.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes the phosphatase 2A catalytic subunit. Protein phosphatase 2A is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. This gene encodes an alpha isoform of the catalytic subunit. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	5516
Other Names	HJS3; NEDLBA; PP2Ac; PP2CA; PP2Calpha; RP-C; PP2Abeta; PP2CB; PP2A; D8Ertd766e; Pp2a1; Pp2a2; PP2AB_HUMAN; PPP2CB; PP2A-beta; 3.1.3.16; PP2AA_HUMAN; PPP2CA; PP2A-alpha; Replication protein C (RP-C); PP2AB_MOUSE; PP2AA_MOUSE; PP2AB_RAT; PP2AA_RAT; protein phosphatase 2 catalytic subunit alpha; protein phosphatase 2 (formerly 2A), catalytic subunit, alpha isoform; protein phosphatase 2, catalytic subunit, alpha isozyme; protein phosphatase 2A catalytic subunit, alpha isoform
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	PPP2CB (HGNC:9300)
Function	Catalytic subunit of protein phosphatase 2A (PP2A), a serine/threonine phosphatase involved in the regulation of a wide variety of enzymes, signal transduction pathways, and cellular events (Probable). PP2A can modulate the activity of phosphorylase B kinase, casein kinase 2, mitogen-stimulated S6 kinase, and MAP-2 kinase. Part of the striatin-interacting phosphatase and kinase (STRIPAK) complexes. STRIPAK complexes have critical roles in protein (de)phosphorylation and are regulators of multiple signaling pathways including Hippo, MAPK, nuclear receptor and cytoskeleton remodeling. Different types of STRIPAK complexes are involved in a variety of biological processes such as cell growth, differentiation, apoptosis, metabolism and immune regulation (PubMed:18782753).
Cellular Location	Cytoplasm. Nucleus. Chromosome, centromere. Cytoplasm, cytoskeleton, spindle pole. Note=In prometaphase cells, but not in anaphase cells, localizes at centromeres. During mitosis, also found at spindle poles

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.