

NLE1 Rabbit pAb

Catalog # AP96817

Product Information

Application	WB
Primary Accession	Q9NVX2
Other Accession	Q9NVX2
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53320
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NLE1
Epitope Specificity	351-450/485
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the NLE1/RSA4 family. Contains 8 WD repeats.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The Notch signaling pathway is an evolutionary conserved system that is involved in intracellular communication. Notch receptors play an important role in development and cell-fate decisions. Notchless is a loss-of-function mutant allele that encodes for protein NLE1 (notchless homolog 1). NLE1 is a 485 amino acid WD40-repeat protein that binds to the cytoplasmic domain of Notch, regulating its signaling activity in <i>Drosophila melanogaster</i> and in mice. Deletion of the NLE1 gene in mice during the early stages of development results in embryonic death, while gene deletion in the late stages of development leads to activation of a caspase-3-dependent apoptotic pathway. In plants, NLE1 is crucial for normal cellular growth and development. Under-expression during shoot proliferation causes pleiotropic defects such as delayed flowering and abnormal organ maturation. It may also play a role in 60S ribosomal subunit biogenesis in yeast. NLE1 contains eight WD40 domains and produces one isoform due to alternative splicing.

Additional Information

Gene ID	54475
Other Names	HUSSY7; NLE; Rsa4; I11Jus1; I11Jus4; RGD1308301; NLE1_HUMAN; NLE1; NLE1_MOUSE; notchless homolog 1; notchless homolog (<i>Drosophila</i>)
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	NLE1
Function	Plays a role in regulating Notch activity. Plays a role in regulating the expression of CDKN1A and several members of the Wnt pathway, probably via its effects on Notch activity. Required during embryogenesis for inner mass cell survival (By similarity).
Cellular Location	Nucleus, nucleolus {ECO:0000269 PubMed:12429849, ECO:0000269 Ref.5}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.