

DDX1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9683a

Product Information

Application	WB, E
Primary Accession	Q92499
Other Accession	A2VD92 , Q641Y8 , Q91VR5 , Q4R7L5 , Q0IIK5
Reactivity	Human
Predicted	Bovine, Monkey, Mouse, Rat, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB24517
Calculated MW	82432
Antigen Region	156-185

Additional Information

Gene ID	1653
Other Names	ATP-dependent RNA helicase DDX1, DEAD box protein 1, DEAD box protein retinoblastoma, DBP-RB, DDX1
Target/Specificity	This DDX1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 156-185 amino acids from the N-terminal region of human DDX1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DDX1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DDX1
Function	Acts as an ATP-dependent RNA helicase, able to unwind both RNA-RNA and RNA-DNA duplexes. Possesses 5' single-stranded RNA overhang nuclease

activity. Possesses ATPase activity on various RNA, but not DNA polynucleotides. May play a role in RNA clearance at DNA double-strand breaks (DSBs), thereby facilitating the template-guided repair of transcriptionally active regions of the genome. Together with RELA, acts as a coactivator to enhance NF-kappa-B-mediated transcriptional activation. Acts as a positive transcriptional regulator of cyclin CCND2 expression. Binds to the cyclin CCND2 promoter region. Associates with chromatin at the NF-kappa-B promoter region via association with RELA. Binds to poly(A) RNA. May be involved in 3'-end cleavage and polyadenylation of pre-mRNAs. Component of the tRNA-splicing ligase complex required to facilitate the enzymatic turnover of catalytic subunit RTCB: together with archease (ZBTB8OS), acts by facilitating the guanylation of RTCB, a key intermediate step in tRNA ligation (PubMed:[24870230](#)). Component of a multi-helicase-TICAM1 complex that acts as a cytoplasmic sensor of viral double-stranded RNA (dsRNA) and plays a role in the activation of a cascade of antiviral responses including the induction of pro-inflammatory cytokines via the adapter molecule TICAM1. Specifically binds (via helicase ATP-binding domain) on both short and long poly(I:C) dsRNA (By similarity).

Cellular Location

Nucleus. Cytoplasm. Cytoplasmic granule. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q91VR5}. Mitochondrion {ECO:0000250|UniProtKB:Q91VR5}. Note=Localized with MBNL1, TIAL1 and YBX1 in stress granules upon stress. Localized with CSTF2 in cleavage bodies. Forms large aggregates called DDX1 bodies. Relocalized into multiple foci (IR-induced foci or IRIF) after IR treatment, a process that depends on the presence of chromosomal DNA and/or RNA-DNA duplexes. Relocalized at sites of DNA double-strand breaks (DSBs) in an ATM-dependent manner after IR treatment. Colocalized with RELA in the nucleus upon TNF-alpha induction. Enters into the nucleus in case of active transcription while it accumulates in cytosol when transcription level is low (PubMed:24608264). Colocalizes in the cytosol with DDX21, DHX36 and TICAM1. Colocalizes in the mitochondria with TICAM1 and poly(I:C) RNA ligand. The multi-helicase-TICAM1 complex may translocate to the mitochondria upon poly(I:C) stimulation (By similarity) {ECO:0000250|UniProtKB:Q91VR5, ECO:0000269|PubMed:24608264}

Tissue Location

Highest levels of transcription in 2 retinoblastoma cell lines and in tissues of neuroectodermal origin including the retina, brain, and spinal cord.


Background

DDX1 box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein of unknown function.

References

- # Medland, S.E., et al. *Am. J. Hum. Genet.* 85(5):750-755(2009)
- # Trynka, G., et al. *Gut* 58(8):1078-1083(2009)
- # Tanaka, K., et al. *Oncogene* 28(21):2142-2151(2009)
- # Maggi, L.B. Jr., et al. *Mol. Cell. Biol.* 28(23):7050-7065(2008)
- # Li, L., et al. *Mol. Cell. Biol.* 28(20):6413-6425(2008)
- # Rikova, K., et al. *Cell* 131(6):1190-1203(2007)
- # Golembowski, S., et al. *Immunobiology* 201(5):631-644(2000)

Images



Western blot analysis of DDX1 Antibody (N-term) (Cat. #AP9683a) in K562 cell line lysates (35ug/lane). DDX1 (arrow) was detected using the purified Pab.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.