

# TLR4 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP9685c

## Product Information

---

<b>Application</b>	WB, FC, E
<b>Primary Accession</b>	<a href="#">O00206</a>
<b>Other Accession</b>	<a href="#">Q68Y56</a> , <a href="#">Q9WV82</a> , <a href="#">Q9GL65</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Bovine, Hamster, Pig
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB24528
<b>Calculated MW</b>	95680
<b>Antigen Region</b>	669-698

## Additional Information

---

<b>Gene ID</b>	7099
<b>Other Names</b>	Toll-like receptor 4, hToll, CD284, TLR4
<b>Target/Specificity</b>	This TLR4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 669-698 amino acids from the Central region of human TLR4.
<b>Dilution</b>	WB~~1:1000 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	TLR4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	TLR4
<b>Function</b>	Transmembrane receptor that functions as a pattern recognition receptor recognizing pathogen- and damage-associated molecular patterns (PAMPs and DAMPs) to induce innate immune responses via downstream signaling

pathways (PubMed:[10835634](#), PubMed:[15809303](#), PubMed:[16622205](#), PubMed:[17292937](#), PubMed:[17478729](#), PubMed:[20037584](#), PubMed:[20711192](#), PubMed:[23880187](#), PubMed:[27022195](#), PubMed:[29038465](#), PubMed:[17803912](#)). At the plasma membrane, cooperates with LY96 to mediate the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed:[27022195](#)). Also involved in LPS-independent inflammatory responses triggered by free fatty acids, such as palmitate, and Ni(2+) (PubMed:[20711192](#)). Mechanistically, acts via MYD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:[10835634](#), PubMed:[21393102](#), PubMed:[27022195](#), PubMed:[36945827](#), PubMed:[9237759](#)). Alternatively, CD14- mediated TLR4 internalization via endocytosis is associated with the initiation of a MYD88-independent signaling via the TICAM1-TBK1-IRF3 axis leading to type I interferon production (PubMed:[14517278](#)). In addition to the secretion of proinflammatory cytokines, initiates the activation of NLRP3 inflammasome and formation of a positive feedback loop between autophagy and NF-kappa-B signaling cascade (PubMed:[32894580](#)). In complex with TLR6, promotes inflammation in monocytes/macrophages by associating with TLR6 and the receptor CD86 (PubMed:[23880187](#)). Upon ligand binding, such as oxLDL or amyloid-beta 42, the TLR4:TLR6 complex is internalized and triggers inflammatory response, leading to NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway (PubMed:[23880187](#)). In myeloid dendritic cells, vesicular stomatitis virus glycoprotein G but not LPS promotes the activation of IRF7, leading to type I IFN production in a CD14- dependent manner (PubMed:[15265881](#), PubMed:[23880187](#)). Required for the migration-promoting effects of ZG16B/PAUF on pancreatic cancer cells.

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Early endosome. Cell projection, ruffle {ECO:0000250|UniProtKB:Q9QUK6}. Note=Upon complex formation with CD36 and TLR6, internalized through dynamin-dependent endocytosis (PubMed:20037584). Colocalizes with RFTN1 at cell membrane and then together with RFTN1 moves to endosomes, upon lipopolysaccharide stimulation. Co-localizes with ZG16B/PAUF at the cell membrane of pancreatic cancer cells (PubMed:36232715)

#### Tissue Location

Highly expressed in placenta, spleen and peripheral blood leukocytes (PubMed:9237759, PubMed:9435236). Detected in monocytes, macrophages, dendritic cells and several types of T-cells (PubMed:27022195, PubMed:9237759). Expressed in pancreatic cancer cells but not in normal pancreatic cells (at protein level) (PubMed:36232715).

## Background


TLR4 is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor is most abundantly expressed in placenta, and in myelomonocytic subpopulation of the leukocytes. It has been implicated in signal transduction events induced by lipopolysaccharide (LPS) found in most gram-negative bacteria.

## References


- # Sam-Agudu, N.A., et al. Am. J. Trop. Med. Hyg. 82(4):548-555(2010)
- # Palomino-Morales, R.J., et al. Arthritis Res. Ther. 12 (2), R51 (2010)
- # Rigoli, L., et al. Anticancer Res. 30(2):513-517(2010)

## Images

---



Western blot analysis of TLR4 Antibody (Center) (Cat. #AP9685c) in HL-60 cell line lysates (35ug/lane). TLR4 (arrow) was detected using the purified Pab.



TLR4 Antibody (Center) (Cat. #AP9685c) flow cytometric analysis of U251 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

## Citations

---

- [Activation of the porcine alveolar macrophages via toll-like receptor 4/NF-κB mediated pathway provides a mechanism of resistin leading to inflammation.](#)
- [Activation of porcine alveolar macrophages by Actinobacillus pleuropneumoniae lipopolysaccharide via the TLR4/NF-κB mediated pathway.](#)
- [Migration ability and Toll-like receptor expression of human mesenchymal stem cells improves significantly after three-dimensional culture.](#)
- [Zhikang Capsule ameliorates dextran sodium sulfate-induced colitis by inhibition of inflammation, apoptosis, oxidative stress and MyD88-dependent TLR4 signaling pathway.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.