

hnRNP DL Rabbit pAb

Catalog # AP96895

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q14979
Other Accession	Q14979
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46438
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human hnRNP DL
Epitope Specificity	101-200/420
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 2 RRM (RNA recognition motif) domains.
Post-translational modifications	Dimethylation of Arg-408 is probably of the asymmetric type.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two RRM domains that bind to RNAs. Three alternatively spliced transcript variants have been described for this gene. One of the variants is probably not translated because the transcript is a candidate for nonsense-mediated mRNA decay. The protein isoforms encoded by this gene are similar to its family member HNRPD. [provided by RefSeq, May 2011]

Additional Information

Gene ID	9987
Other Names	HNRPD; hnRNP DL; HNRNP; JKTBP; JKTBP2; LGMD1G; LGMD3; laAUF1; D5Ert650e; D5Wsu145e; Hnrnpd1; hnRNP-DL; HNRDL_CHICK; HNRNPDL; hnRNP D-like; HNRDL_HUMAN; AU-rich element RNA-binding factor; JKT41-binding protein; Protein laAUF1; HNRDL_MOUSE; HNRDL_RAT; heterogeneous nuclear ribonucleoprotein D like; limb girdle muscular dystrophy 1G (autosomal dominant)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	HNRNPDL
Synonyms	HNRPDL, JKTBP
Function	Acts as a transcriptional regulator. Promotes transcription repression. Promotes transcription activation in differentiated myotubes (By similarity). Binds to double- and single-stranded DNA sequences. Binds to the transcription suppressor CATR sequence of the COX5B promoter (By similarity). Binds with high affinity to RNA molecules that contain AU-rich elements (AREs) found within the 3'-UTR of many proto-oncogenes and cytokine mRNAs. Binds both to nuclear and cytoplasmic poly(A) mRNAs. Binds to poly(G) and poly(A), but not to poly(U) or poly(C) RNA homopolymers. Binds to the 5'-ACUAGC-3' RNA consensus sequence.
Cellular Location	Nucleus. Cytoplasm. Note=Shuttles between the nucleus and the cytoplasm in a TNPO1-dependent manner.
Tissue Location	Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon and leukocytes. Expressed in myeloid leukemia, gastric adenocarcinoma, cervical carcinoma, hepatoma, fibrosarcoma, colon adenocarcinoma, epidermoid carcinoma, osteosarcoma and urinary bladder carcinoma cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.