

KRT73 Rabbit pAb

Catalog # AP96898

Product Information

Application	WB
Primary Accession	Q86Y46
Other Accession	Q86Y46
Reactivity	Human
Predicted	Rat, Mouse, Rabbit, Dog, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	58923
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KRT73
Epitope Specificity	301-400/540
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the intermediate filament family.
SUBUNIT	Heterotetramer of two type I and two type II keratins.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into epithelial keratins and hair keratins. This gene encodes a protein that is expressed in the inner root sheath of hair follicles. The type II keratins are clustered in a region of chromosome 12q13.[provided by RefSeq, Jun 2009]

Additional Information

Gene ID	319101
Other Names	CK-73; IRT6IRS3; K6IRS3; K73; KRT6IRS3; K2C73_HUMAN; KRT73; Cytokeratin-73 (CK-73); Keratin-73 (K73); Type II inner root sheath-specific keratin-K6irs3; Type-II keratin Kb36; KB36; keratin 73; keratin 73, type II
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	KRT73
-------------	-------

Synonyms	K6IRS3, KB36, KRT6IRS3
Function	Has a role in hair formation. Specific component of keratin intermediate filaments in the inner root sheath (IRS) of the hair follicle (Probable).
Tissue Location	Highly expressed in hair follicles from scalp. In hair, it is specifically present in the inner root sheath (IRS) of the hair follicle. Present in the IRS cuticle, but not in Henle or Huxley layers of the IRS. In the IRS cuticle, it is expressed between the lowermost bulb region of the cuticle and the region where Henle cells undergo abrupt terminal differentiation. Detected up to the uppermost cortex region where cuticle cells terminally differentiate (at protein level).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.