

HPR Rabbit pAb

Catalog # AP96961

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P00739
Other Accession	P00739
Host	Rabbit
Clonality	Polyclonal
Calculated MW	39030
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HPR
Epitope Specificity	101-200/348
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the peptidase S1 family. Contains 1 peptidase S1 domain. Contains 1 Sushi (CCP/SCR) domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	HPR is a ubiquitous transcription factor required for a diverse set of processes. It is a component of the CCR4 complex involved in the control of gene expression.

Additional Information

Gene ID	3250
Other Names	A-259H10.2; HP; HPTR_HUMAN; HPR; haptoglobin-related protein
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	HPR
Function	Primate-specific plasma protein associated with apolipoprotein L-I (apoL-I)-containing high-density lipoprotein (HDL). This HDL particle, termed trypanosome lytic factor-1 (TLF-1), mediates human innate immune protection against many species of African trypanosomes. Binds hemoglobin

with high affinity and may contribute to the clearance of cell-free hemoglobin to allow hepatic recycling of heme iron.

Cellular Location

Secreted. Note=Secreted into blood plasma and associated with subtypes of high density lipoproteins (HDL).

Tissue Location

In adult liver the amount of HPR mRNA is at the lower limit of detection, therefore the extent of its expression is at most less than 1000-fold that of the HP1F gene. No HPR mRNA can be detected in fetal liver. Expressed in Hep-G2 and leukemia MOLT-4 cell lines.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.