

SOX18 Rabbit pAb

Catalog # AP96969

Product Information

Application	WB
Primary Accession	P35713
Other Accession	P35713
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40891
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SOX18
Epitope Specificity	101-200/384
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 HMG box DNA-binding domain. Contains 1 Sox C-terminal domain.
DISEASE	Defects in SOX18 are the cause of hypotrichosis-lymphedema-telangiectasia syndrome (HLTS) [MIM:607823].
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. The encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins. This protein plays a role in hair, blood vessel, and lymphatic vessel development. Mutations in this gene have been associated with recessive and dominant forms of hypotrichosis-lymphedema-telangiectasia. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	54345
Other Names	HLTRS; HLTS; Ra; Ragl; SOX18_HUMAN; SOX18; SOX18_MOUSE; Sox-18; SRY-box transcription factor 18; SRY (sex determining region Y)-box 18; SRY-box 18
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SOX18
Function	Transcriptional activator that binds to the consensus sequence 5'-AACAAAG-3' in the promoter of target genes and plays an essential role in embryonic cardiovascular development and lymphangiogenesis. Activates transcription of PROX1 and other genes coding for lymphatic endothelial markers. Plays an essential role in triggering the differentiation of lymph vessels, but is not required for the maintenance of differentiated lymphatic endothelial cells. Plays an important role in postnatal angiogenesis, where it is functionally redundant with SOX17. Interaction with MEF2C enhances transcriptional activation. Besides, required for normal hair development.
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00267}.
Tissue Location	Detected in heart, lung, placenta, skeletal muscle, liver, kidney, spleen, prostate, ovary, msosmall intestine and colon

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.