

phospho-IKB epsilon (Ser22) Rabbit pAb

Catalog # AP97062

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	O00221
Other Accession	O00221
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52864
Physical State	Liquid
Immunogen	KLH conjugated synthesised phosphopeptide derived from human IKB epsilon around the phosphorylation site of Ser22
Epitope Specificity	LR(p-S)LR
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the NF-kappa-B inhibitor family. Contains 6 ANK repeats.
SUBUNIT	Interacts with RELA, REL, NFKB1 nuclear factor NF-kappa-B p50 subunit and NFKB2 nuclear factor NF-kappa-B p52 subunit.
Post-translational modifications	Serine phosphorylated; followed by proteasome-dependent degradation.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	NFKB1 or NFKB2 is bound to REL, RELA, or RELB to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, or NFKBIB), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine). For some genes, activation requires NFKB interaction with other transcription factors, such as STAT, AP1 (JUN), and NFAT.

Additional Information

Gene ID	4794
Other Names	IKBE; IKBE_HUMAN; NFKBIE; NF-kappa-BIE; I-kappa-B-epsilon (IkB-E IkB-epsilon IkappaBepsilon); NFKB inhibitor epsilon; nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, epsilon; NF-kappa-B inhibitor epsilon; NFKBIE IKB epsilon (phospho-S22); p-IKB epsilon; phospho-IKB epsilon; p-IkBe; NFKBIE IKB epsilon (phospho-Ser22)
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500

Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.
----------------	---

Protein Information

Name	NFKBIE
Synonyms	IKBE
Function	Sequesters NF-kappa-B transcription factor complexes in the cytoplasm, thereby inhibiting their activity (PubMed: 9315679). Sequestered complexes include NFKB1-RELA (p50-p65) and NFKB1-REL (p50- c-Rel) complexes (PubMed: 9135156 , PubMed: 9315679). Limits B-cell activation in response to pathogens, and also plays an important role in B-cell development (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Highly expressed in spleen, testis and lung, followed by kidney, pancreas, heart, placenta and brain. Also expressed in granulocytes and macrophages

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.