

IKB epsilon Rabbit pAb

Catalog # AP97112

Product Information

Application	WB
Primary Accession	O00221
Other Accession	O00221
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52864
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human IKB epsilon
Epitope Specificity	165-270/500
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the NF-kappa-B inhibitor family. Contains 6 ANK repeats.
SUBUNIT	Interacts with RELA, REL, NFKB1 nuclear factor NF-kappa-B p50 subunit and NFKB2 nuclear factor NF-kappa-B p52 subunit.
Post-translational modifications	Serine phosphorylated; followed by proteasome-dependent degradation.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	NFKB1 or NFKB2 is bound to REL, RELA, or RELB to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, or NFKBIB), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine). For some genes, activation requires NFKB interaction with other transcription factors, such as STAT, AP1 (JUN), and NFAT.

Additional Information

Gene ID	4794
Other Names	IKBE; IKBE_HUMAN; NFKBIE; NF-kappa-BIE; I-kappa-B-epsilon (IkB-E IkB-epsilon IkappaBepsilon); NFKB inhibitor epsilon; nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, epsilon; NF-kappa-B inhibitor epsilon
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	NFKBIE
Synonyms	IKBE
Function	Sequesters NF-kappa-B transcription factor complexes in the cytoplasm, thereby inhibiting their activity (PubMed: 9315679). Sequestered complexes include NFKB1-RELA (p50-p65) and NFKB1-REL (p50- c-Rel) complexes (PubMed: 9135156 , PubMed: 9315679). Limits B-cell activation in response to pathogens, and also plays an important role in B-cell development (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Highly expressed in spleen, testis and lung, followed by kidney, pancreas, heart, placenta and brain. Also expressed in granulocytes and macrophages

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.