

ACC Recombinant Rabbit mAb

Catalog # AP97145

Product Information

Application	WB, IHC-P, IHC-F, IF, IP, E
Primary Accession	O00763
Other Accession	O00763
Host	Rabbit
Clonality	Recombinant
Calculated MW	276541
Isotype	IgG
Purity	affinity purified by Protein A

Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Background Descriptions	Acetyl-CoA carboxylase (ACC) is a complex multifunctional enzyme system. ACC is a biotin-containing enzyme which catalyzes the carboxylation of acetyl-CoA to malonyl-CoA, the rate-limiting step in fatty acid synthesis. ACC-beta is thought to control fatty acid oxidation by means of the ability of malonyl-CoA to inhibit carnitine-palmitoyl-CoA transferase I, the rate-limiting step in fatty acid uptake and oxidation by mitochondria. ACC-beta may be involved in the regulation of fatty acid oxidation, rather than fatty acid biosynthesis. There is evidence for the presence of two ACC-beta isoforms. [provided by RefSeq, Jul 2008],

Additional Information

Gene ID	32
Other Names	ACC, Acetyl-CoA carboxylase 2;ACC-beta;Acetyl-CoA carboxylase 1;ACC1;ACACA;ACAC;ACCA;ACC-alpha;ACACB, Acetyl CoA carboxylase 2, ACC beta, ACC2, ACCB, AcetylCoA carboxylase 2, ACCbeta, ACC β , ACC β ,
Dilution	IHC-P=1:50-200,IHC-F=1:50-200,WB=1:500-2000,IF=1:50-200,ELISA=1:5000-10000,IP=1:50-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ACACB (HGNC:85)
Synonyms	ACC2, ACCB
Function	Mitochondrial enzyme that catalyzes the carboxylation of acetyl-CoA to

malonyl-CoA and plays a central role in fatty acid metabolism (PubMed:[16854592](#), PubMed:[19236960](#), PubMed:[19900410](#), PubMed:[20457939](#), PubMed:[20952656](#), PubMed:[26976583](#)). Catalyzes a 2 steps reaction starting with the ATP-dependent carboxylation of the biotin carried by the biotin carboxyl carrier (BCC) domain followed by the transfer of the carboxyl group from carboxylated biotin to acetyl-CoA (PubMed:[19236960](#), PubMed:[20457939](#), PubMed:[20952656](#), PubMed:[26976583](#)). Through the production of malonyl-CoA that allosterically inhibits carnitine palmitoyltransferase 1 at the mitochondria, negatively regulates fatty acid oxidation (By similarity). Together with its cytosolic isozyme ACACA, which is involved in de novo fatty acid biosynthesis, promotes lipid storage (By similarity).

Cellular Location

Mitochondrion.

Tissue Location

Widely expressed with highest levels in heart, skeletal muscle, liver, adipose tissue, mammary gland, adrenal gland and colon (PubMed:9099716). Isoform 3 is expressed in skeletal muscle, adipose tissue and liver (at protein level) (PubMed:19190759). Isoform 3 is detected at high levels in adipose tissue with lower levels in heart, liver, skeletal muscle and testis (PubMed:19190759)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.