

Macro H2A.2 Rabbit pAb

Catalog # AP97187

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9P0M6
Other Accession	Q9P0M6
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40058
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Macro H2A.2
Epitope Specificity	1-100/372
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Contains 1 histone H2A domain. Contains 1 Macro domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Additional Information

Gene ID	55506
Other Names	H2AFY2; RGD1561371; H2AW_HUMAN; MACROH2A2; Histone macroH2A2; mH2A2; H2AW_MOUSE; H2afy3; macroH2A.2 histone; H2A histone family, member Y2; H2A histone family member Y2
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	MACROH2A2 (HGNC:14453)
Function	Variant histone H2A which replaces conventional H2A in a subset of nucleosomes where it represses transcription. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and

nucleosome remodeling. May be involved in stable X chromosome inactivation.

Cellular Location

Nucleus. Chromosome. Note=Enriched in inactive X chromosome chromatin (PubMed:11331621, PubMed:11262398) and in senescence- associated heterochromatin (PubMed:15621527)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.