

CXCL2/GRO Beta Rabbit pAb

Catalog # AP97201

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P19875
Other Accession	P19875
Reactivity	Rat, Human, Mouse
Predicted	Pig, Rabbit, Horse, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	11389
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MIP2
Epitope Specificity	51-107/107
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the intercrine alpha (chemokine CxC) family.
Post-translational modifications	The N-terminal processed form GRO-beta(5-73) is produced by proteolytic cleavage after secretion from bone marrow stromal cells.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	GRO beta is a member of the CXC, or chemokine class. It contains the ELR domain immediately preceding the first cysteine residue near the amino terminus. Other chemokines in this group include IL8, GRO alpha/beta/gamma, mouse KC, ENA78, GCP2, PBP/CTAPIII/beta TG/NAP2. These chemokines act primarily on neutrophils as chemoattractants and activators, including neutrophil degradation with release of myeloperoxidase and other enzymes. GRO beta was originally identified as a heparin-binding protein secreted from a murine macrophage cell line in response to endotoxin stimulation. GRO beta is an approximately 8 kDa polypeptide of 73 amino acids. The precursor form of GRO beta consists of 100 amino acids. To generate the mature GRO beta, the precursor cleaves its amino terminal 27 amino acids. GRO beta shows 60% amino acid homology to human GRO alpha and GRO gamma.

Additional Information

Gene ID	2920
Other Names	CINC-2a; GRO2; GROb; MGSA-b; MIP-2a; MIP2; MIP2A; SCYB2; MIP-2; Scyb; CXCL2_HUMAN; CXCL2; Growth-regulated protein beta (Gro-beta); Macrophage inflammatory protein 2-alpha (MIP2-alpha); CXCL2_MOUSE; Macrophage inflammatory protein 2 (MIP2); CXCL2_RAT; Cytokine-induced neutrophil chemoattractant 3 (CINC-3); Cinc3; C-X-C motif chemokine ligand 2;

GRO2 oncogene; chemokine (C-X-C motif) ligand 2

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=0.2 ug/test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	CXCL2
Synonyms	GRO2, GROB, MIP2A, SCYB2
Function	Produced by activated monocytes and neutrophils and expressed at sites of inflammation. Hematopoietic chemokine, which, in vitro, suppresses hematopoietic progenitor cell proliferation. GRO-beta(5-73) shows a highly enhanced hematopoietic activity.
Cellular Location	Secreted.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.