

CRTR1 Rabbit pAb

Catalog # AP97298

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	Q9NZI6
Other Accession	Q9NZI6
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54627
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human CRTR1
Epitope Specificity	181-280/479
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the grh/CP2 family. CP2 subfamily.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The long terminal repeat binding protein-1 (LBP-1) mammalian family of transcription factors are proteins that stimulate transcription in HeLa cells from the major late promoter of simian virus 40 in vitro. The two related human LBP-1 genes are TF2L1, which encodes the alternatively spliced transcripts LBP-1a and LBP-1b, and TF2L2, which encodes LBP-1c and LBP-1d. LBP-9, also designated transcription factor CP2-like 1 (TF2L1), is a protein of 479 amino acids. LBP-9 shows 83% sequence homology with LBP-1b and acts as a suppressor factor inhibiting the stimulation effect of LBP-1b. LBP-9 may regulate the P450scc reporter activity through the -155/-131 element.

Additional Information

Gene ID	29842
Other Names	CRTR1; LBP-9; LBP9; 1810030F05Rik; 4932442M07Rik; Cp2l1; Crtr-1; D930018N21Rik; Tcfcp2l1; TF2L1_HUMAN; TF2L1; CP2-related transcriptional repressor 1 (CRTR-1); Transcription factor LBP-9; TF2L1_MOUSE; transcription factor CP2 like 1; transcription factor CP2-like 1
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TFCP2L1
Synonyms	CRTR1, LBP9
Function	Transcription factor that facilitates establishment and maintenance of pluripotency in embryonic stem cells (ESCs) (PubMed:25215486, PubMed:26906118). With KLF2, acts as the major effector of self-renewal that mediates induction of pluripotency downstream of LIF/STAT3 and Wnt/beta-catenin signaling (By similarity). Required for normal duct development in the salivary gland and kidney (By similarity). Coordinates the development of the kidney collecting ducts intercalated (IC) and principal (PC) cells, which regulate acid- base and salt-water homeostasis, respectively (By similarity). Regulates the expression of IC genes including subunits B1 and D2 of the V-ATPase complex, OXGR1, CA12, SLC4A1, AQP6 and IC-specific transcription factor FOXI1 (By similarity). Also regulates the expression of JAG1 and subsequent notch signaling in the collecting duct (By similarity). JAG1 initiates notch signaling in PCs but inhibits notch signaling in ICs (By similarity). Acts as a transcriptional suppressor that may suppress UBP1-mediated transcriptional activation (By similarity). Modulates the placental expression of CYP11A1 (PubMed:10644752).
Cellular Location	Nucleus.
Tissue Location	Highly expressed in placental JEG-3 cells and very low levels of expression in non-steroidogenic cells. No expression was seen in adrenal NCI-H295A cells or in adrenal tissue

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.