

DUSP5 Rabbit pAb

Catalog # AP97316

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q16690
Other Accession	Q16690
Reactivity	Rat, Mouse
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42047
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DUSP5
Epitope Specificity	221-320/384
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily. Contains 1 rhodanese domain. Contains 1 tyrosine-protein phosphatase domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. This gene product inactivates ERK1, is expressed in a variety of tissues with the highest levels in pancreas and brain, and is localized in the nucleus. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	1847
Other Names	DUSP; HVH3; Gm337; Cpg21; DUS5_HUMAN; DUSP5; Dual specificity protein phosphatase hVH3; 3.1.3.16; 3.1.3.48; VH3; DUS5_RAT; MAP-kinase phosphatase CPG21; dual specificity phosphatase 5
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	DUSP5
Synonyms	VH3
Function	Dual specificity protein phosphatase; active with phosphotyrosine, phosphoserine and phosphothreonine residues. The highest relative activity is toward ERK1.
Cellular Location	Nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.