

VPS25 Rabbit pAb

Catalog # AP97427

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q9BRG1
Other Accession	Q9BRG1
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20748
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human VPS25
Epitope Specificity	21-120/176
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the VPS25 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a protein that is a subunit of the endosomal sorting complex required for transport II (ESCRT-II). This protein complex functions in sorting of ubiquitinated membrane proteins during endocytosis. A pseudogene of this gene is present on chromosome 1. [provided by RefSeq, Jul 2013]

Additional Information

Gene ID	84313
Other Names	DERP9; EAP20; FAP20; 1110020N13Rik; D11Wsu68e; VPS25_HUMAN; VPS25; hVps25; Dermal papilla-derived protein 9; ELL-associated protein of 20 kDa; ESCRT-II complex subunit VPS25; VPS25_MOUSE; vacuolar protein sorting 25 homolog; vacuolar protein sorting 25 (yeast); vacuolar protein sorting 25 homolog (S. cerevisiae)
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	VPS25
-------------	-------

Synonyms	DERP9, EAP20
Function	Component of the ESCRT-II complex (endosomal sorting complex required for transport II), which is required for multivesicular body (MVB) formation and sorting of endosomal cargo proteins into MVBs. The MVB pathway mediates delivery of transmembrane proteins into the lumen of the lysosome for degradation. The ESCRT-II complex is probably involved in the recruitment of the ESCRT-III complex. The ESCRT-II complex may also play a role in transcription regulation, possibly via its interaction with ELL. The ESCRT-II complex may be involved in facilitating the budding of certain RNA viruses.
Cellular Location	Cytoplasm. Endosome membrane. Nucleus, nucleoplasm. Note=Distributes diffusely throughout the cytoplasm and nucleoplasm, but exhibits a punctate distribution on coexpression with CHMP6
Tissue Location	Expressed at the mRNA level in kidney, liver, pancreas, and placenta. Lower levels of expression are found in heart, skeletal muscle, brain and lung.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.