

ORC2 Rabbit pAb

Catalog # AP97435

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q13416
Other Accession	Q13416
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65972
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ORC2
Epitope Specificity	201-300/577
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the ORC2 family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The origin recognition complex (ORC) is a highly conserved six subunits protein complex essential for the initiation of the DNA replication in eukaryotic cells. Studies in yeast demonstrated that ORC binds specifically to origins of replication and serves as a platform for the assembly of additional initiation factors such as Cdc6 and Mcm proteins. The protein encoded by this gene is a subunit of the ORC complex. This protein forms a core complex with ORC3, -4, and -5. It also interacts with CDC45 and MCM10, which are proteins known to be important for the initiation of DNA replication. This protein has been demonstrated to specifically associate with the origin of replication of Epstein-Barr virus in human cells, and is thought to be required for DNA replication from viral origin of replication. Alternatively spliced transcript variants have been found, one of which is a nonsense-mediated mRNA decay candidate. [provided by RefSeq, Oct 2010]

Additional Information

Gene ID	4999
Other Names	ORC2L; ORC2_BOVIN; ORC2; ORC2_HUMAN; ORC2_MOUSE; ORC2_RAT; origin recognition complex subunit 2; origin recognition complex, subunit 2 (yeast homolog)-like; origin recognition complex, subunit 2-like (yeast); origin recognition complex, subunit 2 homolog (yeast); origin recognition complex, subunit 2
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ORC2
Synonyms	ORC2L
Function	Component of the origin recognition complex (ORC) that binds origins of replication. DNA-binding is ATP-dependent. The specific DNA sequences that define origins of replication have not been identified yet. ORC is required to assemble the pre-replication complex necessary to initiate DNA replication. Binds histone H3 and H4 trimethylation marks H3K9me3, H3K20me3 and H4K27me3. Stabilizes LRWD1, by protecting it from ubiquitin-mediated proteasomal degradation. Also stabilizes ORC3.
Cellular Location	Nucleus.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.