

SLITRK4 Rabbit pAb

Catalog # AP97461

Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q8IW52
Other Accession	Q8IW52
Host	Rabbit
Clonality	Polyclonal
Calculated MW	94331
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SLITRK4
Epitope Specificity	251-350/837
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the SLITRK family. Contains 12 LRR (leucine-rich) repeats. Contains 2 LRRCT domains. Contains 1 LRRNT domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	SLITRK family proteins are integral membrane proteins that have a C-terminal domain that is partially similar to TRK neurotrophin receptor proteins and two leucine-rich repeat (LRR) domains that are similar to those of SLIT proteins. SLITRK4 (SLIT and NTRK-like protein 4) is a 837 amino acid single-pass type I membrane protein that contains 18 LRR (leucine-rich) repeats and is expressed in neural tissues, specifically in the thalamus, hypothalamus, subventricular zone, CA3 region of the hippocampus and cortical plate. SLITRK4 may be upregulated in some astrocytic brain tumors such as glioblastomas, astrocytomas and primitive neuroectodermal tumors. As compared with its family member SLITRK2, SLITRK4 only weakly suppresses neurite outgrowth. A study using genome-wide transcriptional profiling suggested that the gene encoding SLITRK4, as well as the ARL5B and PLA2G7 genes, may be involved in the pathogenesis of preeclampsia.

Additional Information

Gene ID	139065
Other Names	6430513G13; D930039I09Rik; SLIK4_HUMAN; SLITRK4; SLIK4_MOUSE; SLIT and NTRK like family member 4; SLIT and NTRK-like family, member 4
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-500,ELISA=1:500 0-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SLITRK4
Function	It is involved in synaptogenesis and promotes synapse differentiation (PubMed: 27812321). Suppresses neurite outgrowth (By similarity).
Cellular Location	Membrane; Single-pass type I membrane protein. Cell membrane
Tissue Location	Expressed in the cerebral cortex of the brain and at higher levels in some astrocytic brain tumors such as astrocytomas, glioblastomas and primitive neuroectodermal tumors

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.